
Technical paper n°7/2014

**Pre-scoping document for the Continental/Pannonian
Steppic/ Black Sea regions
(1st part: Core document)**

Mora Aronsson, Marita Arvela, Jérôme Bailly-Maitre,
Laura Patricia Gavilan, Dominique Richard, Annika Sohlman

10 December 2014

Authors' affiliation:

Mora Aronsson, Swedish Agricultural University, SLU (SE)
Marita Arvela, previously from Muséum national d'Histoire Naturelle, MNHN (FR)
Jérôme Bailly-Maitre, Muséum national d'Histoire naturelle, MNHN (FR)
Laura Patricia Gavilan, Muséum national d'Histoire naturelle, MNHN (FR)
Dominique Richard, Muséum national d'Histoire naturelle, MNHN (FR)
Annika Sohlman, Swedish Agricultural University, SLU (SE)

EEA project manager:

Ivone Pereira Martins

ETC/BD production support:

Muriel Vincent, Muséum national d'Histoire naturelle (FR)

Context:

The Topic Centre has prepared this Technical paper in collaboration with the European Environment Agency (EEA) under its 2014 work programme as a contribution to the EEA's work on Nature legislation, implementation, reporting and assessments.

Citation:

Please cite this report as

Aronsson, M., Arvela, M., Bailly-Maitre, J., Gavilan, L-P., Richard, D. and A. Sohlman, 2014. Pre-scoping document for the Continental/ Pannonian/ Steppic and Black Sea regions (1st part Core document). ETC/BD report to the EEA, 27 pp + annexes.

©ETC/BD 2014

ETC/BD Technical paper N°7/2014

European Topic Centre on Biological Diversity

c/o Muséum national d'Histoire naturelle

57 rue Cuvier

75231 Paris cedex, France

Phone: + 33 1 40 79 38 70

E-mail: etc.biodiversity@mnhn.fr

Website: <http://bd.eionet.europa.eu/>

Contents

| | |
|---|-----------|
| 1. Background | 4 |
| 2. General information on the Continental, Pannonian, Steppic and Black Sea regions..... | 5 |
| 3. Use of Article 17 reporting data on conservation status | 12 |
| 4. Identifying habitat types & species for the Continental, Pannonian, Steppic and Black Sea Seminar – preparatory work..... | 16 |
| 4.1 Data and method used for the analysis and ranking of species and habitat types and habitat groups | 16 |
| 4.1.1 Data used..... | 16 |
| 4.1.2 Methods used..... | 16 |
| 4.2 Results of habitat and species ranking | 21 |
| 4.2.1 Continental region | 21 |
| 4.2.2 Pannonian region | 23 |
| 5. Final selection of habitat-types by the Steering Committee..... | 25 |
| 6. Introduction to background information sheets for the 59 selected habitat-types..... | 27 |
| 7. Appendix 1. List of species and habitats types of the Continental region.... | 28 |
| 8. Appendix 2. List of species and habitats types of the Pannonian region... | 46 |
| 9. Appendix 3. List of species and habitats types occurring only in Bulgaria and/or Romania..... | 56 |

1. Background

The new Natura 2000 Seminars at the biogeographical level aim to exchange and analyse information on measures necessary to achieving favourable conservation status of species and habitats of Community interest, with special attention to the management and coherence of the Natura 2000 network. The seminars involve Member States, key user groups, NGOs and independent experts.

It is important to keep in mind that the Natura 2000 Seminars under discussion are a new process and should not be confused with the biogeographical seminars examining the Member State proposals for SCIs which started in the late 1990s.

The draft Terms of Reference for the new process of Natura 2000 seminars dated 8.4.2011 identifies a pre-scoping phase with the following preparatory work (page 3 and 4):

- *Background work to identify relevant criteria to focus further analysis and discussions (e.g. focus on species and habitats related to ecosystems that are of special importance or under particular threat for a biogeographical region, focus on the most threatened species and habitats or focus on species and habitats for which response to measures is likely, focus on habitats that provide important ecosystem services, including in relation to climate change mitigation and adaptation, not to select species and habitats present only in one Member State or where already at favourable conservation status).*
- *Identifying the species and habitat types considered to be priorities for discussion at a seminar, using existing data from the biogeographical region and the Article 17 reporting process, also having regard to the nature sub-target of the new EU biodiversity strategy.*
- *A consultation phase with Member States, Commission and stakeholders to agree which criteria to use and to decide on the species and habitat types or clusters of species or habitat types that will finally be selected. The expert group on Natura 2000 management being the most appropriate forum for this consultation.*

NB: The "pre-scoping documents" for Natura 2000 Seminars are developed in two or three stages; at each drafting stage there will be additional information and sections. Draft versions are subsequently expanded and completed to take into account decisions by the Steering Committees of each region.

The document is targeted to serve the discussion and planning of the seminar for the Continental, Pannonian, Steppic and Black Sea regions.

The 1st (draft) pre-scoping document dated 11.12.2013 by the ETC/BD was part of the pre-scoping phase and followed largely the approach developed for the pilot seminar at the Boreal and the seminar processes for the Atlantic, Alpine and Mediterranean and Macaronesian regions. It described the methodology to rank the habitat types and species and provided some additional information based mainly on the Art 17 data from the first reporting round (2001-2006).

Based on this first pre-scoping document but also as a result of discussions among Member States, the Steering Committee of the Continental, Pannonian, Steppic and Black Sea regions selected a list of 59 habitat-types for priority discussions.

The second pre-scoping document follows up on the first pre-scoping document and adds information related to the 59 selected habitat-types, making use this time of results from the second Art 17 reporting round (2007-2012). This document will contribute to the preparation of the seminar background document which will be drafted by the Commission consultant.

2. General information on the Continental, Pannonian, Steppic and Black Sea regions

The European Union has nine biogeographical regions (map 2.1), each with its own characteristic blend of vegetation, climate and geology. Working at the biogeographical level makes it easier to discuss conservation of species and habitat types under similar natural conditions across a suite of countries, irrespective of political and administrative boundaries.

Map 2.1 The biogeographical regions of the European Union





Table 2.1 Natura 2000 sites per biogeographical region


| Biogeographical region | Area of Natura 2000 sites per region (km ²)* | Number of sites |
|------------------------|--|-----------------|
| Alpine | 225378,24 | 1964 |
| Atlantic | 297343,27 | 3668 |
| Black sea | 19535,79 | 57 |
| Boreal | 133328,86 | 6762 |
| Continental | 371991,31 | 9993 |
| Macaronesia | 8845,33 | 263 |
| Mediterranean | 358150,4 | 4269 |
| Pannonic | 36115,015 | 896 |
| Steppic | 18176,18 | 99 |

Source: European Natura 2000 database. *Calculation made from tabular data.

Figure 2.1 Total area of Natura 2000 sites (%) in nine biogeographical regions (Natura 2000 database, end 2013)



Map 2.2 Natura 2000 sites across the Continental, Pannonian, Steppic and Black Sea region



Continental region

The Continental region covers 24 countries and 14 of them are EU Member States: Austria, Belgium, Bulgaria, Croatia, the Czech Republic, Denmark, France, Germany, Italy, Luxembourg, Poland, Romania, Slovakia and Sweden.

The Continental region connects to most other biogeographical regions of Europe. It extends in a central-west band over most of Europe. A relatively narrow fringe of land separates it from the Atlantic Ocean in the west; in the east it reaches as far as the border of Asia, just south of the Ural Mountains. It reaches Denmark and Sweden in the north, Italy and the Balkan Peninsula in the south. The region includes or borders to several European Alpine biogeographical subregions. The region is not entirely contiguous: the Alps act as a barrier, isolating the part of the region on the Apennine Peninsula. The Continental region entirely surrounds the Pannonian region as well as the Carpathian Mountains. The Czech Republic is fully within the region except for a small area in southeast. Luxembourg is wholly within the region.

- The Continental biogeographical region is the second largest biogeographical region in Europe, nearly as big as the Boreal region
- The climate is continental with warm summers and cold winters, especially in the central and eastern parts
- The soils have naturally high fertility and the region is a main crop-producing area, mostly through intensive agriculture, but alternative farming is increasing
- Permanent grasslands are still widespread, but the area is decreasing due to intensification of agriculture and afforestation
- The forested area is increasing. The natural forest cover in much of the region is deciduous, but coniferous forests now dominate in several countries
- Only little remains of natural old forests. Substantial areas of forests with little or no management are still found in the eastern parts. In many areas the forest condition is still seriously affected by long-range air pollution
- The urban areas are among the largest and most extensive in Europe, with a dense and increasing infrastructure leading to continuously increasing fragmentation of habitats
- A large proportion of the population lives in the vicinity of forests. Some new afforestation occurs around big cities for recreational purposes
- Most of Europe's large rivers cross the region; often they are highly regulated and the floodplains drained. Many rivers are interconnected by canals
- Wetlands have been much reduced, but some large remaining bogs and mires still occur in the central and eastern parts
- The number of indigenous species is high, but few are endemic to the region; forests and grasslands contain the largest number of species
- All big carnivores exist in the region, which also hosts a reconstituted population of the largest wild European herbivore, the bison
- This is a most important region for birds, including migratory species

Source: EEA (2003) Europe's biodiversity – biogeographical regions and seas. Biogeographical regions in Europe

For further information on the Continental region, please see:

- European Commission (2009) *Natura 2000 in the Continental region* (characteristics of the region, number of Annex I habitat types and Annex II species compared to other biogeographical regions etc.);
<http://ec.europa.eu/environment/nature/info/pubs/docs/biogeos/Continental.pdf>

- EEA (2003) Europe's biodiversity – biogeographical regions and seas. Biogeographical regions in Europe. The Continental biogeographical region – agriculture, fragmentation and big rivers. http://www.eea.europa.eu/publications/report_2002_0524_154909/biogeographical-regions-in-europe/continental_biogeographical_region.pdf

Pannonian region

The Pannonian region is completely surrounded by mountains. It is enclosed by the Alps in the west and the Dinarics in the south. The Carpathians encircle the north and east. As regards the main features of relief, alluvial plains dominate with sparse isolated low hills in the interior and low mountain ranges along the boundaries. The main feature of the region is the Great Hungarian plain. Other plains include the Danube plain in Slovakia. The hilly landscape west of the Danube includes several small mountain ranges as the Bakony and Mecsek hills in Hungary and Fruska Gora hills in Serbia, Papuk. The northern rim is composed of volcanic mountains (Berecse, Pilis, Cserhát, Bukk and Zemplin hills). The fluvial network is an important feature of the region. The Danube which flows from north to south has numerous tributaries. The Pannonian region is very rich in underground water.

Four EU member states are concerned by the Pannonian region, namely the Czech Republic, Hungaria, Romania and Slovakia.

- The Pannonian region is dominated by the Great Hungarian Plain.
- Former extensive forests are replaced by grasslands and steppes. Sandy grasslands, i. e. the Hungarian Puszta, is now the dominating type of habitat.
- Agriculture, drainage, eutrophication and salinisation are major threats to biodiversity.
- River regulation and effluents in river are imposing threats to biodiversity in water bodies.
- A few large lakes are heavily influenced by eutrophication and tourist activities.

Source: EEA (2003) Europe's biodiversity – biogeographical regions and seas. Biogeographical regions in Europe.

For further information on the Pannonian region, please see:

- European Commission (2005) *Natura 2000 in the Pannonian region* (characteristics of the region, number of Annex I habitat types and Annex II species compared to other biogeographical regions etc.)
<http://ec.europa.eu/environment/nature/info/pubs/docs/biogeos/pannonian.pdf>
- EEA (2003) Europe's biodiversity – biogeographical regions and seas. Biogeographical regions in Europe. The Pannonian region – the remains of the Pannonian Sea.
http://www.eea.europa.eu/publications/report_2002_0524_154909/biogeographical-regions-in-europe/pannonian.pdf

Steppic region

The Steppic region stretches from Bucharest in the west, across the lower section of the floodplain of the Danube, along the north of the Black Sea and the foothills of the Caucasus. It is bordered in the east by the northwest coast of the Caspian Sea and the Ural River. Its boundary in the north is the beginning of the wooded steppe, which is part of the Continental region. It represents the European part of the steppes, a continuous band as far as to the Altai Mountains on the borders of Mongolia. Only Romania shares this region in the EU territory.

- Vast steppic areas with a tree-less vegetation dominated by *Stipa* and other turf grasses growing on black soils
- Heavily influenced by human activities as conversion of the steppe into arable land, regulation of waterways, constructions of dams, salinisation and pollution from mining activities
- The Volga delta offers breeding grounds for many birds species
- Uncontrolled use of biological resources has diminished animal populations, several species are listed as vulnerable or endangered, among them sturgeons

Source: EEA (2003) Europe's biodiversity – biogeographical regions and seas. Biogeographical regions in Europe.

For further information on the Steppic region, please see:

- European Commission (2009) *Natura 2000 in the Steppic region* (characteristics of the region, number of Annex I habitat types and Annex II species compared to other biogeographical regions etc.) <http://ec.europa.eu/environment/nature/info/pubs/docs/biogeos/Steppic.pdf>
- EEA (2003) Europe's biodiversity – biogeographical regions and seas. Biogeographical regions in Europe. The Steppic region – the plains of Europe. http://www.eea.europa.eu/publications/report_2002_0524_154909/biogeographical-regions-in-europe/steppic.pdf

Black Sea Region

Most of the Black Sea region can be found in Turkey and from the EU countries only Romania and Bulgaria share this region.

The region consists of two coastal bands encompassing the southern half of the Black Sea. The western part stretches from the delta of the Danube, through the Dobrouja plateau, across low mountains extending east towards the Bosphorus outlet – 1 400 km long and between 10 and 160 km wide – stretches from east of the Bosphorus over the various mountain ranges along the southern coast of the Black Sea and as far the Caucasus mountains in the east.

- This chapter covers southern shore areas of the Black Sea.
- Climatic transfer from Mediterranean to continental Europe.
- Costal cliffs and the Danube delta hosts a highly diverse fauna.
- Originally mostly covered by forests.
- Agriculture, afforestation, salinisation and coastal erosion are major threats to the biodiversity in region.

Source: EEA (2003) Europe's biodiversity – biogeographical regions and seas. Biogeographical regions in Europe.

For further information on the Black Sea region, please see:

- European Commission (2009) *Natura 2000 in the Black Sea region* (characteristics of the region, number of Annex I habitat types and Annex II species compared to other biogeographical regions etc.) <http://ec.europa.eu/environment/nature/info/pubs/docs/biogeos/Black%20Sea.pdf>
- EEA (2003) Europe's biodiversity – biogeographical regions and seas. Biogeographical regions in Europe. The Black Sea region – shores and delta http://www.eea.europa.eu/publications/report_2002_0524_154909/regional-seas-around-europe/BlackSea.pdf

General:

Natura 2000 Barometer provides figures for the global Natura 2000 sites (SPAs+ SCIs) which have been obtained by GIS analysis, using the electronic spatial boundaries provided by Member States for each of their sites. It is regularly updated in Natura 2000 Newsletter:

http://ec.europa.eu/environment/nature/info/pubs/natura2000nl_en.htm

3. Use of Article 17 reporting data on conservation status

- All Member States are required by the Habitats Directive to monitor habitat types and species of Community interest. Article 17 of the Directive requires that every 6 years Member States prepare reports to be sent to the European Commission on the implementation of the Directive. This includes assessments on the conservation status of the habitat types and species of Community interest, *i.e.* habitats listed in Annex I and species listed in Annex II, IV and V of the Habitats Directive across the whole territory of the Member State concerned (not only within Natura 2000).
- The Article 17 reports prepared by the Member States have three sections; (i) general information about the implementation of the Habitats Directive, (ii) the assessments of conservation status of species and (iii) for habitats. Conservation status was assessed using a standard methodology to facilitate comparisons between Member States and to allow aggregation to give assessments for biogeographical regions. Conservation status is assessed as being either ‘favourable’ (FV), ‘unfavourable-inadequate’ (U1) and ‘unfavourable-bad’ (U2), based on four parameters as defined in Article 1 of the Directive.
- The parameters for habitat types are ‘range’, ‘area covered by the habitat type’, ‘structure and functions’ and ‘future prospects’ and for species they are ‘range’, ‘population’, ‘habitat of species’ and ‘future prospects’. Member States were encouraged to use expert opinions where there was insufficient data to inform judgements. However, where there was great uncertainty it was also possible to report the conservation status as ‘unknown’. The assessments of the four parameters were combined following an agreed method to give an overall assessment of conservation status. The conservation status is assessed separately for each of the biogeographical region occurring in a Member State.
- When the first draft of the pre-scoping document for the Continental, Pannonian, Steppic and Black Sea regions was prepared in December 2013, only results from the first Art 17 reporting round (2001-2006) were available. Therefore, the methodology for ranking priority habitats for discussion, as described in chapter 4 of the present document was based on Art 17 assessments from that period.
- In the meantime, results from the second Art 17 reporting round (2007-2012) have been published (see: <http://bd.eionet.europa.eu/article17/reports2012/>).
- The below figures reflect the conservation status of habitat types and species assessed as part of this second reporting round.
- Figures 3.1 to 3.4 show the percentages of overall assessments of habitat types and species in each class for each MAES ecosystem-type respectively in the Continental, the Pannonian, the Steppic and the Black Sea regions. The number in brackets shows the number of assessments in each group. The statistics are based on the Member State level biogeographical assessments of conservation status. All habitat types and Annex II and IV species are used in the graph.

Figure 3.1 Conservation status of habitat types and species per habitat group in the Continental region (number of assessments in brackets). Overall assessments at MS level (based on Art. 17 2007-2012)




Figure 3.2 Conservation status of habitat types and species per habitat group in the Pannonian region (number of assessments in brackets). Overall assessments at MS level




Figure 3.3 Conservation status of habitat types and species per habitat group in the Black Sea region (number of assessments in brackets). Overall assessments at MS level





Figure 3.4 Conservation status of habitat types and species per habitat group in the Steppic region (number of assessments in brackets). Overall assessments at MS level



Figures 3.5 and 3.6 show respectively the conservation status of habitat types and species among different biogeographical and marine regions.

Figure 3.5 Conservation status of habitat types per biogeographical and marine region





Figure 3.6 Conservation status of species per biogeographical and marine regions



4. Identifying habitat types & species for the Continental, Pannonian, Steppic and Black Sea Seminar – preparatory work

This chapter summarises the work provided by the ETC/BD to help the Steering Committee in narrowing down the selection of habitat types and species during the preparatory stage of the Continental, Pannonian, Steppic and Black Sea Natura 2000 Seminar. The methodology and ranking of species and habitat types is described below. It is based on results from the first Art 17 reporting round (2001-2006), as quality checked assessments from the second Art 17 reporting round (2007-2012) were not yet available at that stage of the process.

Bird species are not covered in the ETC/BD analysis, but birds are part of the Natura 2000 seminars.

4.1 Data and method used for the analysis and ranking of species and habitat types and habitat groups

4.1.1 Data used

One of the aims of the new process of the Natura 2000 seminars at biogeographical level is to assess and discuss how the management of the Natura 2000 network can best contribute to the improvement of the conservation status of the targeted species and habitat types (and status of birds). This is why the main source of information for the identification of the criteria is the Article 17 reports (<http://bd.eionet.europa.eu/article17>) (see also chapter 3). In addition, there are practical reasons for this choice: this information is easily accessible and it is the most recent data at the European level (covering period 2001-2006, EU25 species listed in the Annex II, IV and V and habitat types of Annex I of the Habitats Directive).

Bulgaria and Romania were not yet part of the previous Art 17 reporting round, however these two Member States are taken into account in the calculations for the Continental region and Romania for the Pannonian region as they were having ‘unknown’ status in the analysis to ease their involvement in the selection of the habitat types for these regions. In addition a table has been provided in Appendix 3 to list species and habitat types which occur only in Romanian and Bulgarian parts of the Continental and Black Sea region or the Pannonian or Steppic regions of Romania. However, Croatia has not been taken into account in calculations due to their recent accession to the EU.

The Article 17 data from the Member State level were used in this proposal as the potential measures deriving from this process would be taken at the national level. The Article 17 data quality issues are not repeated here as they are discussed in details in the Article 17 Technical report <http://bd.eionet.europa.eu/article17/chapter2>.

4.1.2 Methods used

For all calculations concerning the species and habitat types listed in the Annex I, II and IV of the Habitats Directive, the Article 17 reporting data of 2001-2006 are used. Annex V species are excluded following the approach of the Pilot Boreal, the Atlantic and Alpine and the Mediterranean and Macaronesian processes. In addition, species and habitat types occurring only in one Member States are excluded from the calculations (following the same approach as in the other seminar processes).

Continental region

There are 157 habitat types listed for the Continental region in Annex I of the Habitats Directive (this work covers 143 habitats).

In total 371 Annex II and IV species of the Continental region are listed in the Habitats Directive and 62 of these species occur only in Bulgaria and Romania (this work covers 221 species).

Pannonian region

There are 56 habitat types listed for the Pannonian region in Annex I of the Habitats Directive (this work covers 54 habitats).

In total 196 Annex II and IV species of the Pannonian region are listed in the Habitats Directive (151 species are covered in this work).

Steppic & Black Sea region (these regions are not part of the ranking, see Appendix 3)

There are 25 habitat types listed for the Steppic region in Annex I of the Habitats Directive. In total 86 Annex II and IV species of the Steppic region are listed in the Habitats Directive.

There are 55 habitat types listed for the Black Sea region in Annex I of the Habitats Directive. In total 91 Annex II and IV species of the Black Sea region are listed in the Habitats Directive.

i) Criteria for prioritisation (Criterion A, B and C)

Given the need to focus on a limited number of issues in the seminar the priority for discussions of habitat types and species was assessed and ranked. Identifying priorities should reflect on one side the conservation ‘urgency/priority’ (unfavourable conservation status and declining trends) and on the other side joint interest of all Member States involved in the seminar (the priority should be given to habitat types and species which occur in most of the countries in the region).

The following criteria based on the Article 17 reporting are proposed to be used for the first step to narrow down the selection of species and habitat types (criteria for prioritisation). There are three criteria A, B and C. This work was developed for the Pilot Seminar.

Criterion A. Number of MS where species/habitat types are present

The proposal is to give a higher weight to species and habitat types which occur in several Member States. Habitat types and species only occurring in one Member State of the biogeographical region or habitat type and species that just have some outliers in the region from e.g. the Continental region are less important to discuss in such a setting than the habitat types and species that are shared by many Member States in the same region and with their main distribution there. As explained above, species and habitat types occurring in one Member State of the Continental (or the Pannonian) region only are left out from the analysis.

If for example a species in Continental region is only present in two Member State it scores only 2 points, but if it is present in all 13 Continental Member States it scores 13 points (as Croatia not part of the calculations). Criterion A has a multiplier effect as shown below under the paragraph ‘Filtering the species and habitat types based on criteria A, B and C’.

Criterion B. Species and habitat types at unfavourable conservation status (U2 & U1 & XX)

The terms of reference for the biogeographical seminars excludes from the discussion species and habitats already at favourable conservation status. This is why species and habitats with favourable

conservation status are not taken into account under criterion B. Species and habitats are allocated a score based on their conservation status in each Member State in the following way:

The habitat/species scores

- 2 points for each Member State in which it has been assessed as Unfavourable-Bad (U2) and
- 1 point if Unfavourable-Inadequate (U1) or Unknown (XX).

and these scores summed up give the overall score.

For instance the habitat ‘Semi-natural dry grasslands’ 6210 in Pannonian region was assessed as follows

$$B = 2(N^{\circ}U2) + N^{\circ}XX = 2*2 + 1*1 = 5$$

- B = score for criterion B
- $N^{\circ}U2, N^{\circ}U1, N^{\circ}XX$ = number of Member States with the conclusion U2, U1, XX.

| Member State | Article 17 evaluation | Score |
|---------------|-----------------------|-------|
| CZ | U2 | 2 |
| HU | U2 | 2 |
| SK | XX | 1 |
| Overall score | | 5 |

This criterion reflects the importance to agree on management for habitat types and species that are far from being at favourable conservation status compared to those ones which are close to favourable status. The higher is the number of Member States with unfavourable conclusions the higher the score. This method works with absolute numbers, it is not sensitive to the percentage of the habitat area or species population having an unfavourable status. For example if the conclusion in two out four Member States is U2 the species has 4 points (considering it is favourable in remaining two Member States). But the species scores 4 points also if the species occur only in two Member States and both have reported U2 conservation status.

On the other hand the score is dependent on the number of Member States where the habitat/or species occurs. The habitats/species present in several Member States have higher probability to get high scores.

Criterion C. Trend information

All species and habitat types that were reported as having a negative trend in the Article 17 reports are taken into account using the following parameters:

| Feature | Trend |
|----------------------|--|
| Species | “Population” |
| | “Habitat for the species” |
| Habitat types | “Area of the habitat type” |
| | <i>Qualifier for “Structure & functions”</i> |

Ideally, the qualifier information (U1-, U2-) could have been used under the parameter “Structure and functions”, however as qualifiers are not used systematically, it is not used under this criteria for the analysis of the Continental and the Pannonian region.

For these parameters each negative trend information (scoring 1) is counted per species or habitat type.

If both parameters for a species are negative in 13 Member States, the score would be 26 points.

$$C = N^\circ \text{ trend1} + N^\circ \text{ trend2}$$

- C = score for criterion C
- $N^\circ \text{ trend1}, N^\circ \text{ trend2}$ = number of Member States where the trend1, trend2 is negative

NB: For the Boreal region (Pilot seminar) the qualifier information for structure and functions (U1-, U2-) was used to support this criterion as this information was available from the Boreal Member States. Information on structure and function is closely linked to potential management needs, so its use can be justified in this context.

In the second reporting round (2007-2012) Member States have used the qualifier in a more systematic way and the information could be used for the future seminars.

ii) Filtering the species and habitat types based on criteria A, B and C and use of the Priority Index

After the scores are given to each habitat type and species according to the criteria A, B and C, the scores are then used to calculate a Priority Index for each species and habitat type. The algorithm for calculation should be understandable and simple and the Steering Committees for the Boreal, Atlantic and Alpine regions agreed to use the sum of scores for unfavourable conservation status and negative trend multiplied by the number of countries where habitat/species is present: $A^*(B+C)$.

The other options for the algorithm are described in the draft pre-scoping document for the Boreal region by the ETC/BD:

<https://circabc.europa.eu/w/browse/b9886a98-1fe2-40f1-a759-053c62748d6c>

iii) Criteria for clustering habitats and species

The first discussions in 2011 on the new Natura 2000 seminars at biogeographical level identified a need to cluster the habitats and species into broader ecosystems. The original clustering of habitat types and species developed by the EEA and the ETC/BD for the EU 2010 Biodiversity Baseline¹ has been used as a basis to group species and habitat types under broad habitat groups for the Boreal, Atlantic and Alpine seminar processes as this was the most recent available grouping covering all Member States and relatively easy to be adjusted for the purposes of these seminars.

In this earlier background work all species and habitat types are allocated to at least one of the thirteen habitat groups (forests, freshwater, wetlands, grasslands, agro-ecosystems, rocks etc.). The ETC adjusted the habitat groups to better reflect the ecological conditions of each region (see e.g. the pre-scoping document for the Atlantic region <https://circabc.europa.eu/w/browse/b9886a98-1fe2-40f1-a759-053c62748d6c>).

However, as part of the MAES² process a new typology of ecosystems has been developed for mapping and assessment. The ETC/BD has taken this into account in the work for the Continental and Pannonian region (in practise no big difference to the work done so far for the other biogeographical regions). The MAES typology was already taken into account in the Mediterranean and Macaronesian pre-scoping document. See the table below.

¹The EU 2010 Biodiversity Baseline provides facts and figures on the state and trends of the different biodiversity and ecosystem components and supports the EU in developing the post-2010 sub-targets and provides factual data for measuring and monitoring progress in the EU from 2011 to 2020 (<http://www.eea.europa.eu/publications/eu-2010-biodiversity-baseline>)

²Mapping and Assessment of Ecosystems and their Services

| MAES typology of ecosystems | | Naming used in pre-scoping work by ETC/BD |
|---------------------------------------|---|---|
| Major ecosystem category (level 1) | Ecosystem type for mapping & assessment (level 2) | |
| Terrestrial | Urban | - |
| | Cropland | - |
| | Grassland | Yes |
| | Woodland and forest | Yes, but we call it 'Forests' |
| | Heathland and scrub | Yes, but we call it 'Heaths & scrubs' |
| | Sparingly vegetated land | Yes (means rock and ice) |
| | Inland wetlands | Yes, but we call it 'Mires and bogs' |
| | Coastal | Yes, although MAES work not completed yet |
| Freshwater | Rivers and lakes | Yes |
| Marine | | Yes, although MAES work not completed yet |

For the ETC/BD analysis the habitat types and species from Art 17 data are grouped under one habitat group only wherever possible and limited to maximum of two habitat groups³ and Annex V species are excluded (as for the other biogeographical regions).

NB: For some species e.g. bats the clustering may be too narrow, because breeding, foraging, resting and wintering habitat can cover more than two habitats.

As a result, the ETC/BD proposes to use eight habitat groups and habitat types and species are attributed to these groups using Article 17 checklist and ETC/BD expert opinion: *Forests, Grasslands, Coastal, Heaths & scrubs, Lakes and rivers, Mires & bogs, Sparingly vegetated land and Marine*

iv) Using the Priority Index to rank habitat groups

To finalise the ranking of different habitat groups, for each group of habitat type and species the cumulative Priority Index was calculated by summing up the index of each habitat and species and then divided it with the number of habitats and species in the group.

We can use the 'Rivers and lakes' group in table 4.1 as an example for how the Priority Index was calculated. First we use the algorithm $A^*(B + C)$. Each species and habitat types linked to 'Rivers and lakes' gets a figure (index) by using this agreed formula. The figures are summed up and divided by the number of species/habitats of listed under 'Rivers and lakes' giving the cumulative Priority Index 80,7 for this habitat group.

³ Please notice that some Annex II/IV species may be missing under relevant habitat group, but normally Member States should be able to pick them up during the process, at latest when habitats and species are selected under discussion.

4.2 Results of habitat and species ranking

4.2.1 Continental region

Ranking of the six habitat groups is shown in table 4.1. The results of this analysis give ‘Mires & bogs’ and ‘Heaths & scrubs’ the highest ranking suggesting that these habitat groups (their habitat types and species) require particular attention in the Continental region.

Table 4.1 The cumulative priority index in CON region using A*(B+C) and the ranking (excluding Annex I habitat types and Annex II & IV species occurring only in one MS)

| Habitat group | Number of species and habitat types | A*(B+C) |
|-------------------------|-------------------------------------|---------|
| Mires & bogs | 84 | 101.8 |
| Heaths & scrubs | 30 | 88.6 |
| Grasslands | 80 | 81.8 |
| Forests | 89 | 81.4 |
| Rivers & lakes | 78 | 80.7 |
| Sparsely vegetated land | 38 | 71.4 |
| Marine | 5 | 54.8 |

Ranking of the habitats and species

Given the need to focus on a limited number of issues in the Natura 2000 seminar, we have used the algorithm A*(B+C) where A = number of MS, B = unfavourable conservation status and C = negative trend, to calculate a Priority Index for each habitat types and species following the steps as described above. We ranked the top 20 habitat types (habitat types with the highest Priority Index) for the Continental region (see below table 4.2. For legend of table 4.2, see Appendix 1).

The results for all species and habitat types are shown in Appendix 1. For information, occurrence of the habitats and species in the Black Sea and Steppic region are also indicated in the tables.

Table 4.2 Top 20 habitat types of the Continental region (several habitat types have same scores)

| N2K code | Description | Priority | CON CS | Positive trends | Criterion A | Criterion B | Criterion C | A*(B+C) | Occurs in BLS | Occurs in STE |
|----------|--|----------|--------|-----------------|-------------|-------------|-------------|---------|---------------|---------------|
| 6410 | Molinia meadows on calcareous, peaty or clayey-silt-laden soils (<i>Molinion caeruleae</i>) | N | U2 | 0 | 13 | 22 | 9 | 403 | X | X |
| 6210 | Semi-natural dry grasslands and scrubland facies on calcareous substrates (<i>Festuco-Brometalia</i>) (* important orchid sites) | N | U2 | 0 | 13 | 19 | 8 | 351 | X | |
| 7230 | Alkaline fens | N | U2 | 0 | 13 | 18 | 8 | 338 | | |
| 6510 | Lowland hay meadows (<i>Alopecurus pratensis</i> , <i>Sanguisorba officinalis</i>) | N | U1 | 0 | 12 | 18 | 8 | 312 | X | X |
| 7140 | Transition mires and quaking bogs | N | U2 | 0 | 13 | 17 | 6 | 299 | | |
| 3130 | Oligotrophic to mesotrophic standing waters with vegetation of the <i>Littorelletea uniflorae</i> and/or of the <i>Isoëto-Nanojuncetea</i> | N | U2 | 0 | 13 | 17 | 5 | 286 | X | X |
| 6230 | Species-rich <i>Nardus</i> grasslands, on siliceous substrates in mountain areas (and submountain areas in Continental Europe) | Y | U2 | 0 | 11 | 18 | 7 | 275 | | |
| 3150 | Natural eutrophic lakes with <i>Magnopotamion</i> or <i>Hydrocharition</i> - type vegetation | N | U2 | 1 | 13 | 17 | 3 | 260 | X | X |
| 4030 | European dry heaths | N | U2 | 2 | 10 | 18 | 6 | 240 | X | |
| 91F0 | Riparian mixed forests of <i>Quercus robur</i> , <i>Ulmus laevis</i> and <i>Ulmus minor</i> , <i>Fraxinus excelsior</i> or <i>Fraxinus angustifolia</i> , along the great rivers (<i>Ulmenion minoris</i>) | N | U2 | 0 | 11 | 17 | 4 | 231 | X | X |
| 7220 | Petrifying springs with tufa formation (<i>Cratoneurion</i>) | Y | U2 | 0 | 13 | 14 | 3 | 221 | X | |
| 5130 | <i>Juniperus communis</i> formations on heaths or calcareous grasslands | N | U2 | 1 | 12 | 14 | 4 | 216 | | |
| 3160 | Natural dystrophic lakes and ponds | N | U2 | 0 | 10 | 15 | 5 | 200 | X | X |
| 91E0 | Alluvial forests with <i>Alnus glutinosa</i> and <i>Fraxinus excelsior</i> (<i>Alno-Padion</i> , <i>Alnion incanae</i> , <i>Salicion albae</i>) | Y | U2 | 1 | 11 | 14 | 4 | 198 | X | |
| 91D0 | Bog woodland | Y | U2 | 1 | 11 | 14 | 4 | 198 | | |
| 3260 | Water courses of plain to montane levels with the <i>Ranunculion fluitantis</i> and <i>Callitricho-Batrachion</i> vegetation | N | U1 | 0 | 13 | 14 | 1 | 195 | X | X |
| 9110 | Luzulo-Fagetum beech forests | N | U2 | 3 | 13 | 13 | 1 | 182 | | |
| 3140 | Hard oligo-mesotrophic waters with benthic vegetation of <i>Chara</i> spp. | N | U2 | 0 | 12 | 14 | 1 | 180 | X | X |
| 7110 | Active raised bogs | Y | U2 | 0 | 9 | 15 | 4 | 171 | | |
| 7150 | Depressions on peat substrates of the <i>Rhynchosporion</i> | N | U2 | 0 | 10 | 14 | 3 | 170 | | |

4.2.2 Pannonian region

Ranking of the six habitat groups is shown in table 4.3. The results of this analysis give ‘Grasslands’ and ‘Heaths and scrubs’ habitat groups highest scores suggesting that these habitat groups (their habitat types and species) require particular attention in the Pannonian region.

Table 4.3 The cumulative priority index in PAN region using A*(B+C) and the ranking (excluding Annex I habitat types and Annex II & IV species occurring only in one MS)

| Habitat group | Number of species and habitat types | A*(B+C) |
|-------------------------|-------------------------------------|---------|
| Grasslands | 64 | 15.1 |
| Heaths & scrubs | 22 | 14.2 |
| Mires & bogs | 46 | 13 |
| Rivers & lakes | 44 | 12.3 |
| Forests | 61 | 11.8 |
| Sparsely vegetated land | 16 | 6.9 |

Ranking of the habitats and species

Given the need to focus on a limited number of issues in the Natura 2000 seminar, we have used the algorithm A*(B+C) where A = number of MS, B = unfavourable conservation status and C = negative trend, to calculate a Priority Index for each habitat types and species following the steps as described above. We ranked the top 22 habitat types (habitat types with the highest Priority Index) for the Pannonian region (see below table 4.4. For legend, see Appendix 1). The results for all species and habitat types are shown in Appendix 2. Occurrence of the habitats and species in the Black Sea and Steppic region are indicated in the tables.

Table 4.4 Top 22 habitat types of the Pannonian region (several habitat types have same score)

| N2K code | Description | Priority | PAN CS | Positive trends | Criterion A | Criterion B | Criterion C | A*(B+C) | Occurs in BLS | Occurs in STE |
|----------|---|----------|--------|-----------------|-------------|-------------|-------------|---------|---------------|---------------|
| 6260 | Pannonic sand steppes | Y | U2 | 0 | 4 | 7 | 4 | 44 | X | X |
| 91F0 | Riparian mixed forests of Quercus robur, Ulmus laevis and Ulmus minor, Fraxinus excelsior or Fraxinus angustifolia, along the great rivers (Ulmenion minoris) | N | U2 | 0 | 4 | 7 | 2 | 36 | X | X |
| 91I0 | Euro-Siberian steppic woods with Quercus spp. | Y | U2 | 0 | 4 | 7 | 2 | 36 | X | X |
| 6250 | Pannonic loess steppic grasslands | Y | U2 | 0 | 4 | 6 | 3 | 36 | | |
| 6410 | Molinia meadows on calcareous, peaty or clayey-silt-laden soils (Molinion caeruleae) | N | U2 | 0 | 4 | 6 | 2 | 32 | X | X |
| 6440 | Alluvial meadows of river valleys of the Cnidion dubii | N | U2 | 0 | 4 | 6 | 2 | 32 | X | X |
| 40A0 | Subcontinental peri-Pannonic scrub | Y | U2 | 1 | 4 | 6 | 2 | 32 | X | |
| 1530 | Pannonic salt steppes and salt marshes | Y | U2 | 0 | 4 | 5 | 2 | 28 | X | X |
| 6430 | Hydrophilous tall herb fringe communities of plains and of the montane to alpine levels | N | U2 | 0 | 4 | 6 | 1 | 28 | X | X |
| 6510 | Lowland hay meadows (Alopecurus pratensis, Sanguisorba officinalis) | N | U2 | 1 | 4 | 6 | 1 | 28 | X | X |
| 3260 | Water courses of plain to montane levels with the Ranunculion fluitantis and Callitricho-Batrachion vegetation | N | U2 | 0 | 4 | 6 | 1 | 28 | X | X |
| 91E0 | Alluvial forests with Alnus glutinosa and Fraxinus excelsior (Alno-Padion, Alnion incanae, Salicion albae) | Y | U2 | 0 | 3 | 6 | 2 | 24 | X | |
| 6240 | Sub-Pannonic steppic grasslands | Y | U2 | 0 | 3 | 5 | 3 | 24 | X | |
| 7230 | Alkaline fens | N | U2 | 0 | 3 | 5 | 3 | 24 | | |
| 3150 | Natural eutrophic lakes with Magnopotamion or Hydrocharition - type vegetation | N | U1 | 0 | 4 | 5 | 1 | 24 | X | X |
| 3270 | Rivers with muddy banks with Chenopodion rubri p.p. and Bidention p.p. vegetation | N | U1 | 0 | 4 | 6 | 0 | 24 | X | X |
| 6210 | Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia) (* important orchid sites) | N | U2 | 0 | 3 | 5 | 2 | 21 | X | |
| 3130 | Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or of the Isoëto-Nanojuncetea | N | U1 | 0 | 4 | 5 | 0 | 20 | X | X |
| 9180 | Tilio-Acerion forests of slopes, scree and ravines | Y | U2 | 0 | 3 | 5 | 1 | 18 | X | |
| 4030 | European dry heaths | N | U2 | 0 | 3 | 4 | 2 | 18 | X | |
| 3160 | Natural dystrophic lakes and ponds | N | U2 | 0 | 3 | 4 | 2 | 18 | X | X |
| 91M0 | Pannonian-Balkanic turkey oak –sessile oak forests | N | U2 | 0 | 3 | 4 | 2 | 18 | X | X |

5. Final selection of habitat-types by the Steering Committee

As part of the Steering Committee meeting held in Brussels on 5 March 2014 and of a subsequent email consultation which took place between 4 and mid June 2014, Member States and stakeholders provided their feedback on the pre-selection of 20 habitat-types for the Continental region and 22 habitat-types for the Pannonian region proposed by the ETC/BD. This resulted in a final selection of 59 habitat-types which are presented in table 5.1.

Table 5.1 New list of habitat-types selected for priority consideration within the Continental/ Pannonian/ Steppic/ Black Sea regions

| Habitat |
|---|
| 8310 - Caves not open to the public |
| COASTAL² |
| 1310 - Salicornia and other annuals colonizing mud and sand |
| 1150 - Coastal lagoons |
| 1210 - Annual vegetation of stony banks |
| 1130 - Estuaries |
| 2110 - Embryonic shifting dunes |
| 2130 *Fixed coastal dunes with herbaceous vegetation (grey dunes) |
| 2190 - Humid dune slacks |
| 1410 - Mediterranean salt meadows (<i>Juncetalia maritimi</i>) |
| 1240 - Vegetated sea cliffs of the Mediterranean coasts with endemic <i>Limonium</i> spp. |
| WOODLAND AND FOREST |
| 9110 - Luzulo-Fagetum beech forests |
| 91E0 - Alluvial forests with <i>Alnus glutinosa</i> and <i>Fraxinus excelsior</i> (<i>Alno-Padion</i> , <i>Alnion incanae</i> , <i>Salicion albae</i>) |
| 9180 - Tilio-Acerion forests of slopes, screes and ravines |
| 91D0 - Bog woodland |
| 91F0 - Riparian mixed forests of <i>Quercus robur</i> , <i>Ulmus laevis</i> and <i>Ulmus minor</i> , <i>Fraxinus excelsior</i> or <i>Fraxinus angustifolia</i> , along the great rivers (<i>Ulmenion minoris</i>) |
| 9160 - Sub-Atlantic and medio-European oak or oak-hornbeam forests of the <i>Carpinion betuli</i> |
| 9170 - Galio-Carpinetum oak-hornbeam forests |
| 91H0 Pannonic woods with <i>Quercus pubescens</i> |
| 92A0 - <i>Salix alba</i> and <i>Populus alba</i> galleries |
| 91I0 - Euro-Siberian steppic woods with <i>Quercus</i> spp. |
| 91G0 - Pannonic woods with <i>Quercus petraea</i> and <i>Carpinus betulus</i> |
| 91M0 - Pannonic-Balkanic turkey oak –sessile oak forests |
| 91AA - Eastern white oak woods |
| 92D0 - Southern riparian galleries and thickets (<i>Nerio-Tamaricetea</i> and <i>Securinegion tinctoriae</i>) |

| GRASSLAND |
|---|
| 6210 - Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia) (* important orchid sites) |
| 6230 - Species-rich Nardus grasslands, on siliceous substrates in mountain areas (and submountain areas in Continental Europe) |
| 6410 - Molinia meadows on calcareous, peaty or clayey-silt-laden soils (Molinion caeruleae) |
| 6430 - Hydrophilous tall herb fringe communities of plains and of the montane to alpine levels |
| 6110 - Rupicolous calcareous or basophilic grasslands of the Alysso-Sedion albi |
| 6510 - Lowland hay meadows (<i>Alopecurus pratensis</i> , <i>Sanguisorba officinalis</i>) |
| 6520 - Mountain hay meadows |
| 6120 - Xeric and calcarious grasslands |
| 6440 - Alluvial meadows of river valleys of the Cnidion dubii |
| 2330 - Inland dunes with open <i>Corynephorus</i> and <i>Agrostis</i> grasslands |
| 1340* - Inland salt meadows |
| 6240 - Sub-Pannonic steppic grasslands |
| 1530 - Pannonic salt steppes and salt marshes |
| 2340 - Pannonic inland dunes |
| 6250 - Pannonic loess steppic grasslands |
| 6260 - Pannonic sand steppes |
| 6420 - Mediterranean tall humid grasslands of the Molinio-Holoschoenion |
| 62C0 - Ponto-Sarmatic steppes |
| HEATHLAND AND SCRUB |
| 4030 - European dry heaths |
| 5130 - <i>Juniperus communis</i> formations on heaths or calcareous grasslands |
| 40A0 - Subcontinental peri-Pannonic scrub |
| 40C0 - Ponto-Sarmatic deciduous thickets |
| WETLANDS |
| 7140 - Transition mires and quaking bogs |
| 7220 - Petrifying springs with tufa formation (Cratoneurion) |
| 7230 - Alkaline fens |
| 7110 - Active raised bogs |
| 7210 - Calcareous fens with <i>Cladium mariscus</i> and species of the <i>Caricion davallianae</i> |
| 7150 - Depressions on peat substrates of the Rhynchosporion |
| 7120 - Degraded raised bogs still capable of natural regeneration |
| RIVERS AND LAKES |
| 3130 - Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or of the Isoëto-Nanojuncetea |
| 3150 - Natural eutrophic lakes with Magnopotamion or Hydrocharition - type vegetation |
| 3260 - Water courses of plain to montane levels with the Ranunculion fluitantis and Callitricho-Batrachion vegetation |
| 3140 - Hard oligo-mesotrophic waters with benthic vegetation of <i>Chara</i> spp. |
| 3160 - Natural dystrophic lakes and ponds |
| 3270 - Rivers with muddy banks with <i>Chenopodium rubri</i> p.p. and <i>Bidention</i> p.p. Vegetation |

6. Introduction to background information sheets for the 59 selected habitat-types

Information sheets have been prepared for each of the 59 habitat-types selected for the Continental/ Pannonian/ Steppic and Black Sea regions. It includes the following information:

- A definition of the habitat-type;
- An information on the conservation status as reported by Member States for the period 2007-2012;
- The occurrence of the habitats in Natura 2000 sites (SCIs);
- A map showing the distribution of the habitat as reported under Art 17 and the Natura 2000 sites designated for this habitat.

The description of these habitat-types is provided in a separate document (see ETC Technical report n° 2014).

7. Appendix 1. List of species and habitats types of the Continental region

- Different colours are used for different habitat groups. Species/habitat types present in only one MS are excluded.
- H= habitat type, A = amphibian, I = invertebrate, M = mammal, P = plant, R = reptile
- **Prio** = priority habitat type or species
- **I, II and IV** refer to Annexes of the Habitats Directive.
- **CON CS** = Conservation status at the Continental region. Red = unfavourable-bad, amber = unfavourable-inadequate, grey = unknown.
- **Positive trend**: positive trend for population & habitat for species or area of the habitat were used. 0 means that there was not any positive trend reported under the used parameters.
- **Criterion A**= number of MS where species/habitat type is present, **Criterion B** = species/habitat types at U2, U1 or unknown status and **Criterion C**= negative trend and **A*(B+C)** = the agreed algorithm.
- Cells in yellow highlight the data that was used for ranking the habitat types and species.

| Habitat group | N2K code | Taxonomical group | Description | Prio | I | I | I | V | CON CS | Positive trends | Criterion A | Criterion B | Criterion C | A*(B+C) | Occur in BLS | Occur in STE |
|---------------|----------|-------------------|---|------|---|---|---|---|--------|-----------------|-------------|-------------|-------------|---------|--------------|--------------|
| Forests | 91F0 | H | Riparian mixed forests of Quercus robur, Ulmus laevis and Ulmus minor, Fraxinus excelsior or Fraxinus angustifolia, along the great rivers (Ulménion minoris) | N | Y | | | | U2 | 0 | 11 | 17 | 4 | 231 | X | X |
| Forests | 91E0 | H | Alluvial forests with Alnus glutinosa and Fraxinus excelsior (Alno-Padion, Alnion incanae, Salicion albae) | Y | Y | | | | U2 | 1 | 11 | 14 | 4 | 198 | X | |
| Forests | 91D0 | H | Bog woodland | Y | Y | | | | U2 | 1 | 11 | 14 | 4 | 198 | | |
| Forests | 9110 | H | Luzulo-Fagetum beech forests | N | Y | | | | U2 | 3 | 13 | 13 | 1 | 182 | | |

| | | | | | | | | | | | | | | | |
|---------|------|---|--|---|---|--|--|--|----|---|----|----|---|-----|-----|
| | | | | | | | | | | | | | | | |
| Forests | 9170 | H | Galio-Carpinetum oak-hornbeam forests | N | Y | | | | U1 | 1 | 10 | 14 | 1 | 150 | X |
| Forests | 9130 | H | Asperulo-Fagetum beech forests | N | Y | | | | U1 | 2 | 12 | 11 | 1 | 144 | |
| Forests | 9180 | H | Tilio-Acerion forests of slopes, screees and ravines | Y | Y | | | | U1 | 1 | 12 | 11 | 0 | 132 | X |
| Forests | 9160 | H | Sub-Atlantic and medio-European oak or oak-hornbeam forests of the Carpinion betuli | N | Y | | | | U1 | 1 | 9 | 10 | 1 | 99 | |
| Forests | 9190 | H | Old acidophilous oak woods with Quercus robur on sandy plains | N | Y | | | | U2 | 0 | 8 | 11 | 1 | 96 | |
| Forests | 9410 | H | Acidophilous Picea forests of the montane to alpine levels (Vaccinio-Piceetea) | N | Y | | | | U2 | 0 | 8 | 10 | 2 | 96 | |
| Forests | 9150 | H | Medio-European limestone beech forests of the Cephalanthero-Fagion | N | Y | | | | U1 | 1 | 10 | 8 | 0 | 80 | |
| Forests | 9110 | H | Euro-Siberian steppic woods with Quercus spp. | Y | Y | | | | U2 | 0 | 5 | 7 | 2 | 45 | X X |
| Forests | 9260 | H | Castanea sativa woods | N | Y | | | | U2 | 0 | 5 | 5 | 1 | 30 | |
| Forests | 91H0 | H | Pannonic woods with Quercus pubescens | Y | Y | | | | U2 | 0 | 5 | 5 | 1 | 30 | X |
| Forests | 2180 | H | Wooded dunes of the Atlantic, Continental and Boreal region | N | Y | | | | U2 | 0 | 4 | 6 | 1 | 28 | X |
| Forests | 91G0 | H | Pannonic woods with Quercus petraea and Carpinus betulus | Y | Y | | | | U2 | 0 | 4 | 7 | 0 | 28 | X |
| Forests | 92A0 | H | Salix alba and Populus alba galleries | N | Y | | | | U1 | 0 | 4 | 5 | 1 | 24 | X X |
| Forests | 91T0 | H | Central European lichen Scots pine forests | N | Y | | | | U1 | 0 | 3 | 5 | 1 | 18 | |
| Forests | 9120 | H | Atlantic acidophilous beech forests with Ilex and sometimes also Taxus in the shrublayer (Quercion robori-petraeae or Illici-Fagenion) | N | Y | | | | XX | 0 | 4 | 4 | 0 | 16 | |

| | | | | | | | | | | | | | | | |
|---------|------|---|--|---|---|---|---|----|---|----|----|----|-----|---|---|
| | | | | | | | | | | | | | | | |
| Forests | 9530 | H | (Sub-) Mediterranean pine forests with endemic black pines | Y | Y | | | U1 | 0 | 4 | 4 | 0 | 16 | | |
| Forests | 91U0 | H | Sarmatic steppe pine forest | N | Y | | | U2 | 0 | 2 | 3 | 2 | 10 | | |
| Forests | 9140 | H | Medio-European subalpine beech woods with Acer and Rumex arifolius | N | Y | | | U1 | 1 | 3 | 3 | 0 | 9 | | |
| Forests | 91L0 | H | Illyrian oak-hornbeam forests (Erythronio-Carpinion) | N | Y | | | FV | 1 | 3 | 2 | 0 | 6 | | |
| Forests | 9340 | H | Quercus ilex and Quercus rotundifolia forests | N | Y | | | FV | 0 | 2 | 2 | 0 | 4 | | |
| Forests | 9430 | H | Subalpine and montane Pinus uncinata forests (* if on gypsum or limestone) | N | Y | | | U1 | 0 | 2 | 1 | 0 | 2 | | |
| Forests | 91K0 | H | Illyrian Fagus sylvatica forests (Aremonio-Fagion) | N | Y | | | FV | 1 | 2 | 1 | 0 | 2 | | |
| Forests | 1084 | I | Osmoderma eremita | Y | | Y | Y | U2 | 0 | 11 | 19 | 8 | 297 | X | X |
| Forests | 1308 | M | Barbastella barbastellus | N | | Y | Y | U1 | 4 | 13 | 15 | 5 | 260 | X | |
| Forests | 1166 | A | Triturus cristatus | N | | Y | Y | U1 | 4 | 11 | 13 | 10 | 253 | | |
| Forests | 1193 | A | Bombina variegata | N | | Y | Y | U2 | 0 | 10 | 13 | 9 | 220 | X | |
| Forests | 1304 | M | Rhinolophus ferrumequinum | N | | Y | Y | U1 | 3 | 11 | 15 | 4 | 209 | X | X |
| Forests | 1209 | A | Rana dalmatina | N | | | Y | U2 | 5 | 10 | 9 | 9 | 180 | X | |
| Forests | 1323 | M | Myotis bechsteinii | N | | Y | Y | XX | 2 | 12 | 12 | 3 | 180 | X | |
| Forests | 1188 | A | Bombina bombina | N | | Y | Y | U1 | 6 | 9 | 11 | 8 | 171 | X | X |
| Forests | 1341 | M | Muscardinus avellanarius | N | | | Y | XX | 2 | 12 | 8 | 6 | 168 | X | |
| Forests | 1322 | M | Myotis nattereri | N | | | Y | U1 | 1 | 11 | 10 | 5 | 165 | | |
| Forests | 1330 | M | Myotis mystacinus | N | | | Y | U1 | 0 | 12 | 9 | 4 | 156 | X | |
| Forests | 1083 | I | Lucanus cervus | N | | Y | | U1 | 0 | 11 | 11 | 3 | 154 | X | X |
| Forests | 1386 | P | Buxbaumia viridis | N | | Y | | XX | 2 | 9 | 12 | 5 | 153 | | |
| Forests | 1214 | A | Rana arvalis | N | | | Y | U1 | 1 | 8 | 8 | 11 | 152 | | |
| Forests | 1052 | I | Hypodryas maturna | N | | Y | Y | U2 | 1 | 8 | 12 | 6 | 144 | | X |
| Forests | 1324 | M | Myotis myotis | N | | Y | Y | XX | 5 | 11 | 11 | 2 | 143 | X | X |
| Forests | 1088 | I | Cerambyx cerdo | N | | Y | Y | U2 | 0 | 9 | 11 | 4 | 135 | X | X |
| Forests | 1363 | M | Felis silvestris | N | | | Y | U2 | 3 | 9 | 10 | 4 | 126 | X | |

| | | | | | | | | | | | | | | | |
|---------|------|---|------------------------------------|---|--|---|---|----|----|----|----|---|-----|---|---|
| Forests | 1361 | M | <i>Lynx lynx</i> | N | | Y | Y | U2 | 2 | 9 | 12 | 2 | 126 | X | |
| Forests | 1312 | M | <i>Nyctalus noctula</i> | N | | Y | | U1 | 0 | 12 | 8 | 2 | 120 | X | |
| Forests | 1902 | P | <i>Cypripedium calceolus</i> | N | | Y | Y | U1 | 2 | 8 | 9 | 5 | 112 | | |
| Forests | 1331 | M | <i>Nyctalus leisleri</i> | N | | Y | | XX | 0 | 10 | 10 | 1 | 110 | | |
| Forests | 1332 | M | <i>Vespertilio murinus</i> | N | | Y | | XX | 3 | 11 | 10 | 0 | 110 | | |
| Forests | 1381 | P | <i>Dicranum viride</i> | N | | Y | | U1 | 2 | 10 | 8 | 3 | 110 | | |
| Forests | 1317 | M | <i>Pipistrellus nathusii</i> | N | | Y | | U1 | 1 | 12 | 8 | 1 | 108 | X | |
| Forests | 1207 | A | <i>Rana lessonae</i> | N | | Y | | XX | 1 | 7 | 8 | 7 | 105 | | |
| Forests | 1281 | R | <i>Elaphe longissima</i> | N | | Y | | U1 | 2 | 7 | 8 | 7 | 105 | X | |
| Forests | 1326 | M | <i>Plecotus auritus</i> | N | | Y | | U1 | 1 | 11 | 6 | 3 | 99 | | |
| Forests | 1320 | M | <i>Myotis brandtii</i> | N | | Y | | U1 | 0 | 11 | 8 | 0 | 88 | | |
| Forests | 4068 | P | <i>Adenophora liliifolia</i> | N | | Y | Y | U2 | 1 | 5 | 8 | 5 | 65 | | |
| Forests | 1087 | I | <i>Rosalia alpina</i> | Y | | Y | Y | U1 | 2 | 7 | 7 | 2 | 63 | X | |
| Forests | 5009 | M | <i>Pipistrellus pygmaeus</i> | N | | Y | | XX | 2 | 8 | 6 | 1 | 56 | X | |
| Forests | 1352 | M | <i>Canis lupus</i> | Y | | Y | Y | U2 | 5 | 7 | 8 | 0 | 56 | X | X |
| Forests | 1079 | I | <i>Limoniscus violaceus</i> | N | | Y | | U2 | 1 | 5 | 9 | 2 | 55 | | |
| Forests | 1309 | M | <i>Pipistrellus pipistrellus</i> | N | | Y | | FV | 0 | 11 | 5 | 0 | 55 | X | |
| Forests | 4026 | I | <i>Rhysodes sulcatus</i> | N | | Y | | U1 | 0 | 6 | 8 | 1 | 54 | | |
| Forests | 1337 | M | <i>Castor fiber</i> | N | | Y | Y | U1 | 12 | 9 | 5 | 0 | 45 | | |
| Forests | 1305 | M | <i>Rhinolophus euryale</i> | N | | Y | Y | U2 | 0 | 5 | 7 | 2 | 45 | X | |
| Forests | 1167 | A | <i>Triturus carnifex</i> | N | | Y | Y | U1 | 2 | 4 | 5 | 6 | 44 | | |
| Forests | 1078 | I | <i>Callimorpha quadripunctaria</i> | Y | | Y | | FV | 4 | 11 | 4 | 0 | 44 | X | X |
| Forests | 1086 | I | <i>Cucujus cinnaberinus</i> | N | | Y | Y | U1 | 1 | 6 | 5 | 2 | 42 | | |
| Forests | 1354 | M | <i>Ursus arctos</i> | Y | | Y | Y | U2 | 2 | 5 | 6 | 0 | 30 | | |
| Forests | 1191 | A | <i>Alytes obstetricans</i> | N | | Y | | U2 | 0 | 4 | 3 | 4 | 28 | | |
| Forests | 1343 | M | <i>Sicista betulina</i> | N | | Y | | XX | 0 | 5 | 5 | 0 | 25 | | |
| Forests | 1314 | M | <i>Myotis daubentonii</i> | N | | Y | | FV | 4 | 12 | 2 | 0 | 24 | X | |
| Forests | 1936 | I | <i>Anthrenochernes stellae</i> | N | | Y | | U2 | 0 | 4 | 5 | 0 | 20 | | |
| Forests | 5365 | M | <i>Pipistrellus savii</i> | N | | Y | | XX | 1 | 5 | 4 | 0 | 20 | X | |
| Forests | 1993 | A | <i>Triturus dobrogicus</i> | N | | Y | | U2 | 0 | 3 | 4 | 2 | 18 | X | X |
| Forests | 1342 | M | <i>Dryomys nitedula</i> | N | | Y | | XX | 0 | 4 | 3 | 1 | 16 | X | |
| Forests | 1089 | I | <i>Morimus funereus</i> | N | | Y | | FV | 0 | 5 | 3 | 0 | 15 | X | X |
| Forests | 1215 | A | <i>Rana latastei</i> | N | | Y | Y | U1 | 0 | 2 | 2 | 4 | 12 | | |

| | | | | | | | | | | | | | | | |
|------------|------|---|---|---|---|---|----|---|----|----|---|-----|---|---|--|
| Forests | 1328 | M | <i>Nyctalus lasiopterus</i> | N | | Y | XX | 0 | 3 | 4 | 0 | 12 | | | |
| Forests | 2001 | A | <i>Triturus montandoni</i> | N | | Y | U2 | 1 | 3 | 4 | 0 | 12 | | | |
| Forests | 1383 | P | <i>Dichelyma capillaceum</i> | N | | Y | U2 | 0 | 2 | 4 | 1 | 10 | | | |
| Forests | 5003 | M | <i>Myotis alcathoe</i> | N | | Y | XX | 0 | 3 | 3 | 0 | 9 | X | | |
| Forests | 1085 | I | <i>Buprestis splendens</i> | N | | Y | U2 | 0 | 2 | 3 | 1 | 8 | | | |
| Forests | 1243 | R | <i>Algyroides nigropunctatus</i> | N | | Y | U1 | 0 | 2 | 1 | 2 | 6 | | | |
| Forests | 4036 | I | <i>Leptidea morsei</i> | N | | Y | U1 | 0 | 2 | 2 | 1 | 6 | X | | |
| Forests | 1387 | P | <i>Orthotrichum rogeri</i> | N | | Y | XX | 1 | 2 | 2 | 0 | 4 | | | |
| Forests | 4093 | P | <i>Rhododendron luteum</i> | N | | Y | U1 | 1 | 2 | 1 | 1 | 4 | | | |
| Forests | 2097 | P | <i>Paeonia officinalis</i> ssp. <i>banatica</i> | N | | Y | U1 | 1 | 2 | 2 | 0 | 4 | | | |
| Forests | 1927 | I | <i>Stephanopachys</i> <i>substriatus</i> | N | | Y | XX | 0 | 2 | 2 | 0 | 4 | | | |
| Forests | 1939 | P | <i>Agrimonia pilosa</i> | N | | Y | FV | 0 | 2 | 1 | 0 | 2 | X | | |
| Grasslands | 6410 | H | Molinia meadows on calcareous, peaty or clayey-silt-laden soils (Molinion caeruleae) | N | Y | | U2 | 0 | 13 | 22 | 9 | 403 | X | X | |
| Grasslands | 6210 | H | Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco- Brometalia) (* important orchid sites) | N | Y | | U2 | 0 | 13 | 19 | 8 | 351 | X | | |
| Grasslands | 6510 | H | Lowland hay meadows (Alopecurus pratensis, <i>Sanguisorba officinalis</i>) | N | Y | | U1 | 0 | 12 | 18 | 8 | 312 | X | X | |
| Grasslands | 6230 | H | Species-rich <i>Nardus</i> grasslands, on siliceous substrates in mountain areas (and submountain areas in Continental Europe) | Y | Y | | U2 | 0 | 11 | 18 | 7 | 275 | | | |
| Grasslands | 6430 | H | Hydrophilous tall herb fringe communities of plains and of the montane to alpine levels | N | Y | | U1 | 0 | 13 | 12 | 1 | 169 | X | X | |
| Grasslands | 6520 | H | Mountain hay meadows | N | Y | | U1 | 0 | 9 | 13 | 5 | 162 | | | |
| Grasslands | 2330 | H | Inland dunes with open <i>Corynephorus</i> and <i>Agrostis</i> grasslands | N | Y | | U2 | 0 | 8 | 15 | 3 | 144 | | | |

| | | | | | | | | | | | | | | | |
|------------|------|---|--|---|---|--|--|----|---|---|----|---|-----|---|---|
| Grasslands | 6120 | H | Xeric sand calcareous grasslands | Y | Y | | | U2 | 0 | 7 | 12 | 7 | 133 | X | X |
| Grasslands | 1340 | H | Inland salt meadows | Y | Y | | | U2 | 0 | 7 | 10 | 4 | 98 | | |
| Grasslands | 6440 | H | Alluvial meadows of river valleys of the Cnidion dubii | N | Y | | | U2 | 0 | 7 | 11 | 3 | 98 | X | X |
| Grasslands | 2130 | H | Fixed coastal dunes with herbaceous vegetation ("grey dunes") | Y | Y | | | U2 | 0 | 5 | 8 | 3 | 55 | X | X |
| Grasslands | 6130 | H | Calaminarian grasslands of the Violetalia calaminariae | N | Y | | | U1 | 0 | 5 | 7 | 3 | 50 | | |
| Grasslands | 6240 | H | Sub-Pannonic steppic grasslands | Y | Y | | | U1 | 0 | 5 | 6 | 2 | 40 | X | |
| Grasslands | 2120 | H | Shifting dunes along the shoreline with Ammophila arenaria ("white dunes") | N | Y | | | U2 | 0 | 5 | 6 | 1 | 35 | X | |
| Grasslands | 1330 | H | Atlantic salt meadows (Glauco-Puccinellietalia maritimae) | N | Y | | | U2 | 0 | 4 | 7 | 1 | 32 | | |
| Grasslands | 1530 | H | Pannonic salt steppes and salt marshes | Y | Y | | | U2 | 0 | 3 | 4 | 1 | 15 | X | X |
| Grasslands | 6220 | H | Pseudo-steppe with grasses and annuals of the Thero-Brachypodietea | Y | Y | | | FV | 0 | 3 | 3 | 1 | 12 | X | |
| Grasslands | 62A0 | H | Eastern sub-Mediterranean dry grasslands (Scorzoneralia villosae) | N | Y | | | U2 | 0 | 3 | 3 | 1 | 12 | X | |
| Grasslands | 6170 | H | Alpine and subalpine calcareous grasslands | N | Y | | | U1 | 0 | 3 | 2 | 1 | 9 | | |
| Grasslands | 1410 | H | Mediterranean salt meadows (Juncetalia maritimi) | N | Y | | | XX | 0 | 3 | 3 | 0 | 9 | X | X |
| Grasslands | 2340 | H | Pannonic inland dunes | Y | Y | | | U2 | 1 | 2 | 3 | 0 | 6 | | |
| Grasslands | 6250 | H | Pannonic loess steppic grasslands | Y | Y | | | U2 | 0 | 2 | 3 | 0 | 6 | | |
| Grasslands | 6260 | H | Pannonic sand steppes | Y | Y | | | U2 | 0 | 2 | 3 | 0 | 6 | X | X |
| Grasslands | 6190 | H | Rupicolous pannonic grasslands (Stipo-Festucetalia pallentis) | N | Y | | | U1 | 0 | 2 | 2 | 0 | 4 | | |
| Grasslands | 6420 | H | Mediterranean tall humid grasslands of the Molinio-Holoschoenion | N | Y | | | XX | 0 | 2 | 2 | 0 | 4 | X | X |

| | | | | | | | FV | 0 | 3 | 1 | 0 | 3 | | |
|------------|------|---|--|---|---|---|----|----|----|----|----|----|-----|-----|
| | | | | N | Y | | | | | | | | | |
| Grasslands | 6150 | H | Siliceous alpine and boreal grasslands | N | Y | | | | | | | | | |
| Grasslands | 1058 | I | Maculinea arion | N | | Y | | U2 | 0 | 12 | 21 | 16 | 444 | X |
| Grasslands | 1065 | I | Euphydryas aurinia | N | | Y | | U2 | 1 | 13 | 19 | 13 | 416 | X |
| Grasslands | 1261 | R | Lacerta agilis | N | | Y | | U1 | 0 | 11 | 12 | 9 | 231 | X |
| Grasslands | 1056 | I | Parnassius mnemosyne | N | | Y | | U1 | 1 | 9 | 13 | 10 | 207 | X |
| Grasslands | 1197 | A | Pelobates fuscus | N | | Y | | U1 | 2 | 9 | 13 | 9 | 198 | X |
| Grasslands | 1059 | I | Maculinea teleius | N | Y | Y | | U2 | 1 | 8 | 11 | 11 | 176 | |
| Grasslands | 1339 | M | Cricetus cricetus | N | | Y | | U2 | 1 | 8 | 11 | 9 | 160 | |
| Grasslands | 1329 | M | Plecotus austriacus | N | | Y | | XX | 0 | 10 | 10 | 5 | 150 | X |
| Grasslands | 1061 | I | Maculinea nausithous | N | Y | Y | | U2 | 1 | 8 | 11 | 7 | 144 | |
| Grasslands | 1057 | I | Parnassius apollo | N | | Y | | U2 | 0 | 7 | 13 | 7 | 140 | |
| Grasslands | 1321 | M | Myotis emarginatus | N | Y | Y | | U1 | 5 | 11 | 9 | 3 | 132 | X X |
| Grasslands | 1076 | I | Proserpinus proserpina | N | | Y | | XX | 2 | 10 | 10 | 3 | 130 | |
| Grasslands | 1263 | R | Lacerta viridis | N | | Y | | U2 | 0 | 7 | 10 | 8 | 126 | X |
| Grasslands | 1327 | M | Eptesicus serotinus | N | | Y | | XX | 2 | 11 | 7 | 4 | 121 | X |
| Grasslands | 4096 | P | Gladiolus palustris | N | Y | Y | | U1 | 0 | 8 | 9 | 6 | 120 | |
| Grasslands | 1281 | R | Elaphe longissima | N | | Y | | U1 | 2 | 7 | 8 | 7 | 105 | X |
| Grasslands | 1060 | I | Lycaena dispar | N | Y | Y | | U1 | 10 | 11 | 5 | 4 | 99 | X X |
| Grasslands | 1256 | R | Podarcis muralis | N | | Y | | FV | 2 | 9 | 5 | 5 | 90 | X |
| Grasslands | 1307 | M | Myotis blythii | N | Y | Y | | U2 | 0 | 7 | 9 | 3 | 84 | X X |
| Grasslands | 1071 | I | Coenonympha oedippus | N | Y | Y | | U1 | 0 | 5 | 7 | 8 | 75 | X |
| Grasslands | 1310 | M | Miniopterus schreibersii | N | Y | Y | | U2 | 0 | 6 | 9 | 2 | 66 | X X |
| Grasslands | 4038 | I | Lycaena helle | N | Y | Y | | U2 | 0 | 5 | 7 | 5 | 60 | |
| Grasslands | 4030 | I | Colias myrmidone | N | Y | Y | | U1 | 0 | 5 | 7 | 5 | 60 | X |
| Grasslands | 1053 | I | Zerynthia polyxena | N | | Y | | U2 | 1 | 5 | 6 | 5 | 55 | X |
| Grasslands | 335 | M | Spermophilus citellus | N | Y | Y | | U2 | 0 | 5 | 7 | 4 | 55 | X X |
| Grasslands | 1070 | I | Coenonympha hero | N | | Y | | U2 | 0 | 4 | 7 | 5 | 48 | |
| Grasslands | 1437 | P | Thesium ebracteatum | N | Y | Y | | U1 | 0 | 4 | 7 | 2 | 36 | |
| Grasslands | 1617 | P | Angelica palustris | N | Y | Y | | U1 | 1 | 4 | 6 | 3 | 36 | |
| Grasslands | 4067 | P | Echium russicum | N | Y | Y | | U2 | 0 | 4 | 6 | 2 | 32 | X |
| Grasslands | 1279 | R | Elaphe quatuorlineata | N | Y | Y | | U1 | 0 | 4 | 5 | 3 | 32 | X X |
| Grasslands | 4094 | P | Gentianella bohemica | Y | Y | Y | | U2 | 0 | 3 | 5 | 3 | 24 | |
| Grasslands | 2093 | P | Pulsatilla grandis | N | Y | Y | | U1 | 0 | 3 | 4 | 3 | 21 | X |

| | | | | | | | | | | | | | | | | |
|-----------------|------|---|--|---|---|---|----|---|----|----|---|-----|---|---|--|--|
| Grasslands | 1477 | P | Pulsatilla patens | N | Y | Y | U1 | 0 | 4 | 3 | 2 | 20 | | | | |
| Grasslands | 4087 | P | Serratula lycopifolia | Y | Y | Y | U1 | 0 | 4 | 4 | 1 | 20 | | | | |
| Grasslands | 2633 | M | Mustela eversmannii | N | Y | Y | XX | 0 | 4 | 4 | 1 | 20 | X | X | | |
| Grasslands | 1069 | I | Erebia sudetica | N | | Y | U2 | 0 | 3 | 5 | 1 | 18 | | | | |
| Grasslands | 1689 | P | Dracocephalum austriacum | N | Y | Y | U1 | 0 | 3 | 3 | 2 | 15 | | | | |
| Grasslands | 1805 | P | Jurinea cyanoides | Y | Y | Y | U2 | 0 | 2 | 4 | 3 | 14 | | | | |
| Grasslands | 1419 | P | Botrychium simplex | N | Y | Y | FV | 0 | 3 | 3 | 1 | 12 | | | | |
| Grasslands | 1077 | I | Hyles hippophaes | N | | Y | XX | 0 | 3 | 3 | 1 | 12 | X | | | |
| Grasslands | 4013 | I | Carabus hungaricus | N | Y | Y | U2 | 0 | 3 | 4 | 0 | 12 | | | | |
| Grasslands | 1396 | P | Nothothylas orbicularis | N | Y | | U2 | 0 | 2 | 4 | 1 | 10 | | | | |
| Grasslands | 5003 | M | Myotis alcathoe | N | | Y | XX | 0 | 3 | 3 | 0 | 9 | X | | | |
| Grasslands | 4011 | I | Bolbelasmus unicornis | N | Y | Y | XX | 0 | 3 | 3 | 0 | 9 | X | X | | |
| Grasslands | 4091 | P | Crambe tataria | N | Y | Y | U1 | 0 | 3 | 3 | 0 | 9 | | X | | |
| Grasslands | 4035 | I | Gortyna borelii lunata | N | Y | Y | XX | 0 | 3 | 3 | 0 | 9 | | | | |
| Grasslands | 4042 | I | Polyommatus eroides | N | Y | Y | U2 | 0 | 2 | 3 | 1 | 8 | | | | |
| Grasslands | 4104 | P | Himantoglossum adriaticum | N | Y | Y | FV | 0 | 2 | 2 | 1 | 6 | | | | |
| Grasslands | 1298 | R | Vipera ursinii | N | Y | Y | U1 | 0 | 2 | 2 | 1 | 6 | X | | | |
| Grasslands | 4069 | P | Campanula bohemica | Y | Y | Y | U1 | 0 | 2 | 1 | 1 | 4 | | | | |
| Grasslands | 4113 | P | Galium sudeticum | Y | Y | Y | U1 | 2 | 2 | 2 | 0 | 4 | | | | |
| Grasslands | 2021 | M | Sicista subtilis | N | Y | Y | XX | 0 | 2 | 2 | 0 | 4 | | X | | |
| Grasslands | 1547 | P | Genista holopetala | N | Y | Y | U1 | 0 | 2 | 1 | 0 | 2 | | | | |
| Grasslands | 1939 | P | Agrimonia pilosa | N | Y | Y | FV | 0 | 2 | 1 | 0 | 2 | | X | | |
| Heaths & scrubs | 4030 | H | European dry heaths | N | Y | | U2 | 2 | 10 | 18 | 6 | 240 | X | | | |
| Heaths & scrubs | 5130 | H | Juniperus communis formations on heaths or calcareous grasslands | N | Y | | U2 | 1 | 12 | 14 | 4 | 216 | | | | |
| Heaths & scrubs | 4010 | H | Northern Atlantic wet heaths with Erica tetralix | N | Y | | U2 | 0 | 6 | 11 | 6 | 102 | | | | |
| Heaths & scrubs | 2140 | H | Decalcified fixed dunes with Empetrum nigrum | Y | Y | | U1 | 0 | 4 | 6 | 3 | 36 | | | | |
| Heaths & scrubs | 4060 | H | Alpine and Boreal heaths | N | Y | | FV | 0 | 6 | 3 | 1 | 24 | | | | |
| Heaths & scrubs | 2320 | H | Dry sand heaths with Calluna and Empetrum nigrum | N | Y | | U2 | 0 | 3 | 6 | 0 | 18 | | | | |

| | | | | | | | | | | | | | | | |
|-----------------|------|---|---|----------|----------|----------|----------|-----------|----------|-----------|-----------|-----------|------------|----------|----------|
| | | | Bushes with <i>Pinus mugo</i> and <i>Rhododendron hirsutum</i> (<i>Mugo-Rhododendrum hirsuti</i>) | Y | Y | | | U1 | 0 | 4 | 3 | 1 | 16 | | |
| Heaths & scrubs | 4070 | H | Subcontinental peri-Pannonic scrub | Y | Y | | | U1 | 0 | 4 | 3 | 1 | 16 | X | |
| Heaths & scrubs | 2310 | H | Dry sand heaths with <i>Calluna</i> and <i>Genista</i> | N | Y | | | U1 | 0 | 2 | 4 | 1 | 10 | | |
| Heaths & scrubs | 2160 | H | Dunes with <i>Hippophaë rhamnoides</i> | N | Y | | | U1 | 0 | 4 | 2 | 0 | 8 | X | X |
| Heaths & scrubs | 4080 | H | Sub-Arctic <i>Salix</i> spp. scrub | N | Y | | | U1 | 0 | 4 | 2 | 0 | 8 | | |
| Heaths & scrubs | 2250 | H | Coastal dunes with <i>Juniperus</i> spp. | Y | Y | | | U2 | 0 | 2 | 3 | 0 | 6 | | |
| Heaths & scrubs | 5110 | H | Stable xerothermophilous formations with <i>Buxus sempervirens</i> on rock slopes (<i>Berberidion p.p.</i>) | N | Y | | | FV | 0 | 5 | 1 | 0 | 5 | | |
| Heaths & scrubs | 1420 | H | Mediterranean and thermo-Atlantic halophilous scrubs (<i>Sarcocornetea fruticosi</i>) | N | Y | | | XX | 0 | 2 | 1 | 0 | 2 | | |
| Heaths & scrubs | 4090 | H | Endemic oro-Mediterranean heaths with gorse | N | Y | | | FV | 0 | 2 | 1 | 0 | 2 | X | |
| Heaths & scrubs | 5210 | H | Arborescent matorral with <i>Juniperus</i> spp. | N | Y | | | FV | 0 | 2 | 1 | 0 | 2 | X | |
| Heaths & scrubs | 1308 | M | <i>Barbastella barbastellus</i> | N | | Y | Y | U1 | 4 | 13 | 15 | 5 | 260 | X | |
| Heaths & scrubs | 1283 | R | <i>Coronella austriaca</i> | N | | | Y | U1 | 0 | 11 | 11 | 11 | 242 | X | |
| Heaths & scrubs | 1303 | M | <i>Rhinolophus hipposideros</i> | N | | Y | Y | U1 | 5 | 11 | 14 | 7 | 231 | X | X |
| Heaths & scrubs | 1304 | M | <i>Rhinolophus ferrumequinum</i> | N | | Y | Y | U1 | 3 | 11 | 15 | 4 | 209 | X | X |
| Heaths & scrubs | 1056 | I | <i>Parnassius mnemosyne</i> | N | | | Y | U1 | 1 | 9 | 13 | 10 | 207 | X | |
| Heaths & scrubs | 1074 | I | <i>Eriogaster catax</i> | N | | Y | Y | XX | 1 | 10 | 12 | 6 | 180 | | X |
| Heaths & scrubs | 1323 | M | <i>Myotis bechsteinii</i> | N | | Y | Y | XX | 2 | 12 | 12 | 3 | 180 | X | |
| Heaths & scrubs | 1341 | M | <i>Muscardinus avellanarius</i> | N | | | Y | XX | 2 | 12 | 8 | 6 | 168 | X | |
| Heaths & scrubs | 1076 | I | <i>Proserpinus proserpina</i> | N | | | Y | XX | 2 | 10 | 10 | 3 | 130 | | |
| Heaths & scrubs | 1067 | I | <i>Lopinga achine</i> | N | | | Y | U2 | 0 | 8 | 11 | 5 | 128 | | |
| Heaths & scrubs | 1363 | M | <i>Felis silvestris</i> | N | | | Y | U2 | 3 | 9 | 10 | 4 | 126 | X | |
| Heaths & scrubs | 1307 | M | <i>Myotis blythii</i> | N | | Y | Y | U2 | 0 | 7 | 9 | 3 | 84 | X | X |
| Heaths & scrubs | 1313 | M | <i>Eptesicus nilssonii</i> | N | | | Y | U1 | 3 | 8 | 8 | 0 | 64 | | |

| | | | | | | | | | | | | | | |
|-----------------|------|---|--|---|---|---|----|---|----|----|---|-----|---|---|
| Heaths & scrubs | 1352 | M | Canis lupus | Y | Y | Y | U2 | 5 | 7 | 8 | 0 | 56 | X | X |
| Heaths & scrubs | 1053 | I | Zerynthia polyxena | N | | Y | U2 | 1 | 5 | 6 | 5 | 55 | X | |
| Heaths & scrubs | 1070 | I | Coenonympha hero | N | | Y | U2 | 0 | 4 | 7 | 5 | 48 | | |
| Heaths & scrubs | 1078 | I | Callimorpha quadripunctaria | Y | Y | | FV | 4 | 11 | 4 | 0 | 44 | X | X |
| Heaths & scrubs | 4014 | I | Carabus variolosus | N | Y | Y | U1 | 0 | 5 | 5 | 3 | 40 | | |
| Heaths & scrubs | 1050 | I | Saga pedo | N | | Y | U1 | 0 | 5 | 6 | 2 | 40 | | |
| Heaths & scrubs | 1279 | R | Elaphe quatuorlineata | N | Y | Y | U1 | 0 | 4 | 5 | 3 | 32 | X | X |
| Heaths & scrubs | 1316 | M | Myotis capaccinii | N | Y | Y | U2 | 0 | 4 | 5 | 2 | 28 | X | X |
| Heaths & scrubs | 1343 | M | Sicista betulina | N | | Y | XX | 0 | 5 | 5 | 0 | 25 | | |
| Heaths & scrubs | 1342 | M | Dryomys nitedula | N | | Y | XX | 0 | 4 | 3 | 1 | 16 | X | |
| Heaths & scrubs | 1289 | R | Telescopus falax | N | | Y | U2 | 0 | 3 | 4 | 1 | 15 | | |
| Heaths & scrubs | 1077 | I | Hyles hippophaes | N | | Y | XX | 0 | 3 | 3 | 1 | 12 | X | |
| Heaths & scrubs | 1284 | R | Coluber viridiflavus | N | | Y | U1 | 0 | 3 | 2 | 1 | 9 | | |
| Heaths & scrubs | 4011 | I | Bolbelasmus unicornis | N | Y | Y | XX | 0 | 3 | 3 | 0 | 9 | X | X |
| Heaths & scrubs | 1241 | R | Podarcis melisellensis | N | | Y | U1 | 0 | 2 | 2 | 2 | 8 | | |
| Heaths & scrubs | 1243 | R | Algyrodes nigropunctatus | N | | Y | U1 | 0 | 2 | 1 | 2 | 6 | | |
| Heaths & scrubs | 1333 | M | Tadarida teniotis | N | | Y | XX | 0 | 3 | 2 | 0 | 6 | | |
| Marine | 1310 | H | Salicornia and other annuals colonizing mud and sand | N | Y | | U1 | 0 | 9 | 11 | 4 | 135 | X | X |
| Marine | 1150 | H | Coastal lagoons | Y | Y | | U2 | 0 | 6 | 8 | 3 | 66 | X | X |
| Marine | 1130 | H | Estuaries | N | Y | | U2 | 0 | 6 | 5 | 2 | 42 | X | |
| Marine | 1140 | H | Mudflats and sandflats not covered by seawater at low tide | N | Y | | U2 | 0 | 5 | 5 | 0 | 25 | X | |
| Marine | 1320 | H | Spartina swards (Spartinion maritimae) | N | Y | | U1 | 0 | 2 | 2 | 1 | 6 | | |
| Mires & bogs | 7230 | H | Alkaline fens | N | Y | | U2 | 0 | 13 | 18 | 8 | 338 | | |
| Mires & bogs | 7140 | H | Transition mires and quaking bogs | N | Y | | U2 | 0 | 13 | 17 | 6 | 299 | | |
| Mires & bogs | 7220 | H | Petrifying springs with tufa formation (Cratoneurion) | Y | Y | | U2 | 0 | 13 | 14 | 3 | 221 | X | |
| Mires & bogs | 7110 | H | Active raised bogs | Y | Y | | U2 | 0 | 9 | 15 | 4 | 171 | | |

| | | | | | | | | | | | | | | | | |
|--------------|------|---|--|---|---|---|---|---|----|----|----|----|----|-----|---|---|
| Mires & bogs | 7210 | H | Calcareous fens with <i>Cladum mariscus</i> and species of the <i>Caricion davallianae</i> | Y | Y | | | | U1 | 0 | 11 | 10 | 5 | 165 | X | X |
| Mires & bogs | 7120 | H | Degraded raised bogs still capable of natural regeneration | N | Y | | | | U2 | 1 | 8 | 12 | 4 | 128 | | |
| Mires & bogs | 2190 | H | Humid dune slacks | N | Y | | | | U2 | 0 | 5 | 7 | 1 | 40 | X | X |
| Mires & bogs | 2170 | H | Dunes with <i>Salix repens</i> ssp. <i>argentea</i> (<i>Salicion arenariae</i>) | N | Y | | | | XX | 0 | 4 | 4 | 1 | 20 | | |
| Mires & bogs | 1065 | I | <i>Euphydryas aurinia</i> | N | | Y | | | U2 | 1 | 13 | 19 | 13 | 416 | X | |
| Mires & bogs | 1393 | P | <i>Drepanocladus vernicosus</i> | N | | Y | | | U2 | 1 | 11 | 16 | 9 | 275 | | |
| Mires & bogs | 1203 | A | <i>Hyla arborea</i> | N | | | Y | | U1 | 5 | 11 | 13 | 12 | 275 | X | |
| Mires & bogs | 1202 | A | <i>Bufo calamita</i> | N | | | Y | | U2 | 1 | 9 | 17 | 13 | 270 | | |
| Mires & bogs | 1042 | I | <i>Leucorrhinia pectoralis</i> | N | | Y | Y | | U1 | 2 | 11 | 14 | 10 | 264 | | |
| Mires & bogs | 1166 | A | <i>Triturus cristatus</i> | N | | Y | Y | | U1 | 4 | 11 | 13 | 10 | 253 | | |
| Mires & bogs | 1201 | A | <i>Bufo viridis</i> | N | | | Y | | U2 | 2 | 10 | 14 | 10 | 240 | X | |
| Mires & bogs | 1903 | P | <i>Liparis loeselii</i> | N | | Y | Y | | U1 | 3 | 11 | 14 | 7 | 231 | | |
| Mires & bogs | 1303 | M | <i>Rhinolophus hipposideros</i> | N | | Y | Y | | U1 | 5 | 11 | 14 | 7 | 231 | X | X |
| Mires & bogs | 1193 | A | <i>Bombina variegata</i> | N | | Y | Y | | U2 | 0 | 10 | 13 | 9 | 220 | X | |
| Mires & bogs | 1014 | I | <i>Vertigo angustior</i> | N | | Y | | | U1 | 1 | 11 | 11 | 8 | 209 | X | |
| Mires & bogs | 1197 | A | <i>Pelobates fuscus</i> | N | | | Y | | U1 | 2 | 9 | 13 | 9 | 198 | X | |
| Mires & bogs | 1209 | A | <i>Rana dalmatina</i> | N | | | Y | | U2 | 5 | 10 | 9 | 9 | 180 | X | |
| Mires & bogs | 1059 | I | <i>Maculinea teleius</i> | N | | Y | Y | | U2 | 1 | 8 | 11 | 11 | 176 | | |
| Mires & bogs | 1220 | R | <i>Emys orbicularis</i> | N | | Y | Y | | U2 | 2 | 8 | 13 | 9 | 176 | X | X |
| Mires & bogs | 1188 | A | <i>Bombina bombina</i> | N | | Y | Y | | U1 | 6 | 9 | 11 | 8 | 171 | X | X |
| Mires & bogs | 1355 | M | <i>Lutra lutra</i> | N | | Y | Y | | U1 | 10 | 11 | 11 | 3 | 154 | X | X |
| Mires & bogs | 1214 | A | <i>Rana arvalis</i> | N | | | Y | | U1 | 1 | 8 | 8 | 11 | 152 | | |
| Mires & bogs | 1061 | I | <i>Maculinea nausithous</i> | N | | Y | Y | | U2 | 1 | 8 | 11 | 7 | 144 | | |
| Mires & bogs | 1037 | I | <i>Ophiogomphus cecilia</i> | N | | Y | Y | | U1 | 4 | 10 | 9 | 5 | 140 | X | X |
| Mires & bogs | 1016 | I | <i>Vertigo mouliniana</i> | N | | Y | | | U1 | 3 | 10 | 9 | 4 | 130 | X | |
| Mires & bogs | 1093 | I | <i>Austropotamobius torrentium</i> | Y | | Y | | Y | U1 | 2 | 8 | 10 | 6 | 128 | | |
| Mires & bogs | 1312 | M | <i>Nyctalus noctula</i> | N | | | Y | | U1 | 0 | 12 | 8 | 2 | 120 | X | |
| Mires & bogs | 4096 | P | <i>Gladiolus palustris</i> | N | | Y | Y | | U1 | 0 | 8 | 9 | 6 | 120 | | |
| Mires & bogs | 4045 | I | <i>Coenagrion ornatum</i> | N | | Y | | | U1 | 2 | 7 | 10 | 6 | 112 | X | X |

| | | | | | | | | | | | | | | |
|--------------|------|---|-----------------------------|---|--|---|----|----|----|----|---|-----|---|---|
| Mires & bogs | 1207 | A | Rana lessonae | N | | Y | XX | 1 | 7 | 8 | 7 | 105 | | |
| Mires & bogs | 1428 | P | Marsilea quadrifolia | N | | Y | U2 | 1 | 7 | 9 | 6 | 105 | | X |
| Mires & bogs | 1725 | P | Lindernia procumbens | N | | Y | U2 | 0 | 8 | 9 | 4 | 104 | | |
| Mires & bogs | 1318 | M | Myotis dasycneme | N | | Y | U1 | 4 | 9 | 10 | 1 | 99 | | |
| Mires & bogs | 1060 | I | Lycaena dispar | N | | Y | U1 | 10 | 11 | 5 | 4 | 99 | X | X |
| Mires & bogs | 1035 | I | Leucorrhinia caudalis | N | | Y | U2 | 2 | 7 | 9 | 5 | 98 | | |
| Mires & bogs | 1831 | P | Luronium natans | N | | Y | U2 | 1 | 6 | 10 | 6 | 96 | | |
| Mires & bogs | 1048 | I | Aeshna viridis | N | | Y | U2 | 2 | 6 | 8 | 7 | 90 | | |
| Mires & bogs | 1320 | M | Myotis brandtii | N | | Y | U1 | 0 | 11 | 8 | 0 | 88 | | |
| Mires & bogs | 1044 | I | Coenagrion mercuriale | N | | Y | U2 | 1 | 6 | 10 | 4 | 84 | | |
| Mires & bogs | 1071 | I | Coenonympha oedippus | N | | Y | U1 | 0 | 5 | 7 | 8 | 75 | X | |
| Mires & bogs | 1614 | P | Apium repens | N | | Y | U2 | 0 | 5 | 8 | 6 | 70 | | |
| Mires & bogs | 1040 | I | Stylurus flavipes | N | | Y | U1 | 3 | 6 | 7 | 4 | 66 | | |
| Mires & bogs | 1313 | M | Eptesicus nilssonii | N | | Y | U1 | 3 | 8 | 8 | 0 | 64 | | |
| Mires & bogs | 1013 | I | Vertigo geyeri | N | | Y | XX | 1 | 5 | 8 | 4 | 60 | | |
| Mires & bogs | 4038 | I | Lycaena helle | N | | Y | U2 | 0 | 5 | 7 | 5 | 60 | | |
| Mires & bogs | 1292 | R | Natrix tessellata | N | | Y | U1 | 0 | 6 | 7 | 3 | 60 | X | |
| Mires & bogs | 5009 | M | Pipistrellus pygmaeus | N | | Y | XX | 2 | 8 | 6 | 1 | 56 | X | |
| Mires & bogs | 1337 | M | Castor fiber | N | | Y | U1 | 12 | 9 | 5 | 0 | 45 | | |
| Mires & bogs | 1898 | P | Eleocharis carniolica | N | | Y | U1 | 0 | 5 | 6 | 3 | 45 | X | |
| Mires & bogs | 1167 | A | Triturus carnifex | N | | Y | U1 | 2 | 4 | 5 | 6 | 44 | | |
| Mires & bogs | 1038 | I | Leucorrhinia albifrons | N | | Y | U2 | 2 | 5 | 6 | 2 | 40 | | |
| Mires & bogs | 1914 | I | Carabus menetriesi pacholei | Y | | Y | U2 | 1 | 4 | 7 | 3 | 40 | | |
| Mires & bogs | 4014 | I | Carabus variolosus | N | | Y | U1 | 0 | 5 | 5 | 3 | 40 | | |
| Mires & bogs | 1092 | I | Austropotamobius pallipes | N | | Y | U2 | 1 | 4 | 4 | 5 | 36 | | |
| Mires & bogs | 1528 | P | Saxifraga hirculus | N | | Y | U2 | 0 | 3 | 6 | 6 | 36 | | |
| Mires & bogs | 1617 | P | Angelica palustris | N | | Y | U1 | 1 | 4 | 6 | 3 | 36 | | |
| Mires & bogs | 1041 | I | Oxygastra curtisii | N | | Y | U1 | 3 | 5 | 4 | 3 | 35 | | |
| Mires & bogs | 1900 | P | Spiranthes aestivalis | N | | Y | U2 | 0 | 4 | 5 | 3 | 32 | | |
| Mires & bogs | 1191 | A | Alytes obstetricans | N | | Y | U2 | 0 | 4 | 3 | 4 | 28 | | |
| Mires & bogs | 1039 | I | Sympecma braueri | N | | Y | U1 | 1 | 4 | 5 | 2 | 28 | | |
| Mires & bogs | 1316 | M | Myotis capaccinii | N | | Y | U2 | 0 | 4 | 5 | 2 | 28 | X | X |
| Mires & bogs | 1832 | P | Caldesia parnassifolia | N | | Y | U2 | 0 | 3 | 4 | 4 | 24 | | |

| | | | | | | | | | | | | | | |
|----------------|------|---|---|---|---|---|----|---|----|----|---|-----|---|---|
| Mires & bogs | 1516 | P | <i>Aldrovanda vesiculosa</i> | N | Y | Y | U1 | 0 | 4 | 5 | 1 | 24 | | X |
| Mires & bogs | 1993 | A | <i>Triturus dobrogicus</i> | N | Y | | U2 | 0 | 3 | 4 | 2 | 18 | X | X |
| Mires & bogs | 1758 | P | <i>Ligularia sibirica</i> | N | Y | Y | U1 | 0 | 4 | 4 | 0 | 16 | | |
| Mires & bogs | 1217 | R | <i>Testudo hermanni</i> | N | Y | Y | U1 | 0 | 3 | 3 | 2 | 15 | X | X |
| Mires & bogs | 1215 | A | <i>Rana latastei</i> | N | Y | Y | U1 | 0 | 2 | 2 | 4 | 12 | | |
| Mires & bogs | 2001 | A | <i>Triturus montandoni</i> | N | Y | Y | U2 | 1 | 3 | 4 | 0 | 12 | | |
| Mires & bogs | 1383 | P | <i>Dichelyma capillaceum</i> | N | Y | | U2 | 0 | 2 | 4 | 1 | 10 | | |
| Mires & bogs | 1670 | P | <i>Myosotis rehsteineri</i> | N | Y | Y | U2 | 0 | 2 | 3 | 2 | 10 | | |
| Mires & bogs | 1389 | P | <i>Meesia longisetata</i> | N | Y | | U2 | 0 | 2 | 3 | 2 | 10 | | |
| Mires & bogs | 4064 | I | <i>Theodoxus transversalis</i> | N | Y | Y | XX | 0 | 3 | 3 | 0 | 9 | | X |
| Mires & bogs | 1887 | P | <i>Coleanthus subtilis</i> | N | Y | Y | FV | 3 | 3 | 2 | 0 | 6 | | |
| Mires & bogs | 1186 | A | <i>Proteus anguinus</i> | Y | Y | Y | U1 | 0 | 2 | 1 | 2 | 6 | | |
| Mires & bogs | 4046 | I | <i>Cordulegaster heros</i> | N | Y | Y | FV | 0 | 3 | 2 | 0 | 6 | | |
| Mires & bogs | 1714 | P | <i>Euphrasia marchesettii</i> | N | Y | Y | U1 | 0 | 2 | 2 | 0 | 4 | | |
| Mires & bogs | 2217 | P | <i>Pedicularis sudetica</i> | Y | Y | Y | U1 | 1 | 2 | 2 | 0 | 4 | | |
| Mires & bogs | 4093 | P | <i>Rhododendron luteum</i> | N | Y | Y | U1 | 1 | 2 | 1 | 1 | 4 | | |
| Mires & bogs | 1493 | P | <i>Sisymbrium supinum</i> | N | Y | Y | U2 | 0 | 2 | 2 | 0 | 4 | | |
| Mires & bogs | 4019 | I | <i>Leptodirus hochstetii</i> | N | Y | Y | FV | 0 | 2 | 0 | 0 | 0 | | |
| Rivers & lakes | 3130 | H | Oligotrophic to mesotrophic standing waters with vegetation of the <i>Littorelletea uniflorae</i> and/or of the Isoëto-Nanojuncetea | N | Y | | U2 | 0 | 13 | 17 | 5 | 286 | X | X |
| Rivers & lakes | 3150 | H | Natural eutrophic lakes with Magnopotamion or Hydrocharition - type vegetation | N | Y | | U2 | 1 | 13 | 17 | 3 | 260 | X | X |
| Rivers & lakes | 3160 | H | Natural dystrophic lakes and ponds | N | Y | | U2 | 0 | 10 | 15 | 5 | 200 | X | X |
| Rivers & lakes | 3260 | H | Water courses of plain to montane levels with the <i>Ranunculion fluitantis</i> and <i>Callitricho-Batrachion</i> vegetation | N | Y | | U1 | 0 | 13 | 14 | 1 | 195 | X | X |
| Rivers & lakes | 3140 | H | Hard oligo-mesotrophic waters with benthic vegetation of <i>Chara</i> spp. | N | Y | | U2 | 0 | 12 | 14 | 1 | 180 | X | X |

| | | | | | | | | | | | | | | | | |
|----------------|------|---|---|---|---|---|---|---|-----------|----|----|----|----|-----|---|---|
| Rivers & lakes | 3270 | H | Rivers with muddy banks with <i>Chenopodium rubri</i> p.p. and <i>Bidention</i> p.p. vegetation | N | Y | | | | U2 | 0 | 11 | 13 | 2 | 165 | X | X |
| Rivers & lakes | 3220 | H | Alpine rivers and the herbaceous vegetation along their banks | N | Y | | | | U2 | 0 | 7 | 11 | 3 | 98 | | |
| Rivers & lakes | 3240 | H | Alpine rivers and their ligneous vegetation with <i>Salix elaeagnos</i> | N | Y | | | | U2 | 0 | 6 | 9 | 4 | 78 | | X |
| Rivers & lakes | 3110 | H | Oligotrophic waters containing very few minerals of sandy plains (<i>Littorellatalia uniflorae</i>) | N | Y | | | | U1 | 0 | 5 | 8 | 3 | 55 | | |
| Rivers & lakes | 3230 | H | Alpine rivers and their ligneous vegetation with <i>Myricaria germanica</i> | N | Y | | | | U2 | 0 | 4 | 7 | 3 | 40 | | |
| Rivers & lakes | 3180 | H | Turloughs | Y | Y | | | | U1 | 1 | 2 | 1 | 0 | 2 | | |
| Rivers & lakes | 1032 | I | <i>Unio crassus</i> | N | | Y | Y | | U2 | 2 | 12 | 20 | 9 | 348 | X | |
| Rivers & lakes | 1203 | A | <i>Hyla arborea</i> | N | | | Y | | U1 | 5 | 11 | 13 | 12 | 275 | X | |
| Rivers & lakes | 1042 | I | <i>Leucorrhinia pectoralis</i> | N | | Y | Y | | U1 | 2 | 11 | 14 | 10 | 264 | | |
| Rivers & lakes | 1029 | I | <i>Margaritifera margaritifera</i> | N | | Y | | | U2 | 0 | 8 | 16 | 12 | 224 | | |
| Rivers & lakes | 1220 | R | <i>Emys orbicularis</i> | N | | Y | Y | | U2 | 2 | 8 | 13 | 9 | 176 | X | X |
| Rivers & lakes | 1134 | F | <i>Rhodeus sericeus amarus</i> | N | | Y | | | U2 | 5 | 10 | 10 | 7 | 170 | X | X |
| Rivers & lakes | 1145 | F | <i>Misgurnus fossilis</i> | N | | Y | | | U1 | 0 | 9 | 11 | 7 | 162 | X | X |
| Rivers & lakes | 1149 | F | <i>Cobitis taenia</i> | N | | Y | | | U1 | 1 | 12 | 11 | 2 | 156 | X | X |
| Rivers & lakes | 1355 | M | <i>Lutra lutra</i> | N | | Y | Y | | U1 | 10 | 11 | 11 | 3 | 154 | X | X |
| Rivers & lakes | 1037 | I | <i>Ophiogomphus cecilia</i> | N | | Y | Y | | U1 | 4 | 10 | 9 | 5 | 140 | X | X |
| Rivers & lakes | 1082 | I | <i>Graphoderus bilineatus</i> | N | | Y | Y | | FV | 0 | 9 | 12 | 3 | 135 | | X |
| Rivers & lakes | 1093 | I | <i>Austropotamobius torrentium</i> | Y | | Y | | Y | U1 | 2 | 8 | 10 | 6 | 128 | | |
| Rivers & lakes | 1163 | F | <i>Cottus gobio</i> | N | | Y | | | FV | 5 | 12 | 6 | 4 | 120 | | |
| Rivers & lakes | 4056 | I | <i>Anisus vorticulus</i> | N | | Y | Y | | U2 | 0 | 7 | 10 | 6 | 112 | X | X |
| Rivers & lakes | 4045 | I | <i>Coenagrion ornatum</i> | N | | Y | | | U1 | 2 | 7 | 10 | 6 | 112 | X | X |
| Rivers & lakes | 1428 | P | <i>Marsilea quadrifolia</i> | N | | Y | Y | | U2 | 1 | 7 | 9 | 6 | 105 | | X |
| Rivers & lakes | 1096 | F | <i>Lampetra planeri</i> | N | | Y | | | U1 | 6 | 9 | 8 | 3 | 99 | | |
| Rivers & lakes | 1318 | M | <i>Myotis dasycneme</i> | N | | Y | Y | | U1 | 4 | 9 | 10 | 1 | 99 | | |
| Rivers & lakes | 1035 | I | <i>Leucorrhinia caudalis</i> | N | | | Y | | U2 | 2 | 7 | 9 | 5 | 98 | | |

| | | | | | | | | | | | | | | |
|----------------|------|---|----------------------------------|---|---|---|----|---|----|----|---|----|---|---|
| Rivers & lakes | 1831 | P | <i>Luronium natans</i> | N | Y | Y | U2 | 1 | 6 | 10 | 6 | 96 | | |
| Rivers & lakes | 1106 | F | <i>Salmo salar</i> | N | Y | | U2 | 7 | 7 | 12 | 1 | 91 | | |
| Rivers & lakes | 1048 | I | <i>Aeshna viridis</i> | N | | Y | U2 | 2 | 6 | 8 | 7 | 90 | | |
| Rivers & lakes | 1044 | I | <i>Coenagrion mercuriale</i> | N | Y | | U2 | 1 | 6 | 10 | 4 | 84 | | |
| Rivers & lakes | 2484 | F | <i>Eudontomyzon mariae</i> | N | Y | | U1 | 1 | 7 | 10 | 2 | 84 | | |
| Rivers & lakes | 1099 | F | <i>Lampetra fluviatilis</i> | N | Y | Y | U2 | 3 | 6 | 11 | 1 | 72 | | |
| Rivers & lakes | 1130 | F | <i>Aspius aspius</i> | N | Y | Y | U1 | 5 | 9 | 6 | 2 | 72 | X | X |
| Rivers & lakes | 1614 | P | <i>Apium repens</i> | N | Y | Y | U2 | 0 | 5 | 8 | 6 | 70 | | |
| Rivers & lakes | 1040 | I | <i>Stylurus flavipes</i> | N | | Y | U1 | 3 | 6 | 7 | 4 | 66 | | |
| Rivers & lakes | 1138 | F | <i>Barbus meridionalis</i> | N | Y | Y | U2 | 0 | 6 | 7 | 4 | 66 | | |
| Rivers & lakes | 1146 | F | <i>Sabanejewia aurata</i> | N | Y | | U1 | 1 | 6 | 8 | 0 | 48 | X | X |
| Rivers & lakes | 1131 | F | <i>Leuciscus souffia</i> | N | Y | | U1 | 0 | 6 | 5 | 2 | 42 | | |
| Rivers & lakes | 6158 | F | <i>Gobio albipinnatus</i> | N | Y | | U2 | 2 | 5 | 6 | 2 | 40 | X | X |
| Rivers & lakes | 1038 | I | <i>Leucorrhinia albifrons</i> | N | | Y | U2 | 2 | 5 | 6 | 2 | 40 | | |
| Rivers & lakes | 1095 | F | <i>Petromyzon marinus</i> | N | Y | | U1 | 1 | 5 | 7 | 1 | 40 | | |
| Rivers & lakes | 1105 | F | <i>Hucho hucho</i> | N | Y | Y | U2 | 0 | 4 | 5 | 5 | 40 | | |
| Rivers & lakes | 1160 | F | <i>Zingel streber</i> | N | Y | | U2 | 0 | 5 | 6 | 2 | 40 | X | X |
| Rivers & lakes | 1092 | I | <i>Austropotamobius pallipes</i> | N | Y | Y | U2 | 1 | 4 | 4 | 5 | 36 | | |
| Rivers & lakes | 1114 | F | <i>Rutilus pigus</i> | N | Y | Y | U2 | 0 | 4 | 5 | 4 | 36 | | |
| Rivers & lakes | 1041 | I | <i>Oxygastra curtisii</i> | N | Y | Y | U1 | 3 | 5 | 4 | 3 | 35 | | |
| Rivers & lakes | 1081 | I | <i>Dytiscus latissimus</i> | N | Y | Y | U2 | 0 | 4 | 5 | 3 | 32 | | |
| Rivers & lakes | 1159 | F | <i>Zingel zingel</i> | N | Y | Y | U1 | 0 | 5 | 5 | 1 | 30 | X | X |
| Rivers & lakes | 2522 | F | <i>Pelecus cultratus</i> | N | Y | | XX | 0 | 6 | 5 | 0 | 30 | X | X |
| Rivers & lakes | 1102 | F | <i>Alosa alosa</i> | N | Y | Y | U2 | 3 | 4 | 6 | 1 | 28 | | |
| Rivers & lakes | 1103 | F | <i>Alosa fallax</i> | N | Y | Y | U2 | 0 | 4 | 6 | 1 | 28 | X | |
| Rivers & lakes | 1039 | I | <i>Sympecma braueri</i> | N | | Y | U1 | 1 | 4 | 5 | 2 | 28 | | |
| Rivers & lakes | 1122 | F | <i>Gobio uranoscopus</i> | N | Y | | U1 | 0 | 4 | 5 | 2 | 28 | | |
| Rivers & lakes | 2511 | F | <i>Gobio kessleri</i> | N | Y | | U2 | 2 | 5 | 5 | 0 | 25 | | X |
| Rivers & lakes | 1157 | F | <i>Gymnocephalus schraetzeri</i> | N | Y | Y | U1 | 0 | 5 | 5 | 0 | 25 | X | X |
| Rivers & lakes | 1832 | P | <i>Caldesia parnassifolia</i> | N | Y | Y | U2 | 0 | 3 | 4 | 4 | 24 | | |
| Rivers & lakes | 1516 | P | <i>Aldrovanda vesiculosa</i> | N | Y | Y | U1 | 0 | 4 | 5 | 1 | 24 | | X |
| Rivers & lakes | 1314 | M | <i>Myotis daubentonii</i> | N | | Y | FV | 4 | 12 | 2 | 0 | 24 | X | |
| Rivers & lakes | 2555 | F | <i>Gymnocephalus baloni</i> | N | Y | Y | U1 | 0 | 4 | 4 | 0 | 16 | X | X |

| | | | | | | | | | | | | | | |
|-------------------------|------|---|---|---|---|---|----|---|----|----|---|-----|---|---|
| Rivers & lakes | 1217 | R | <i>Testudo hermanni</i> | N | Y | Y | U1 | 0 | 3 | 3 | 2 | 15 | X | X |
| Rivers & lakes | 2011 | F | <i>Umbra krameri</i> | N | Y | | U2 | 0 | 3 | 4 | 1 | 15 | X | X |
| Rivers & lakes | 1115 | F | <i>Chondrostoma genei</i> | N | Y | | U2 | 0 | 2 | 4 | 2 | 12 | | |
| Rivers & lakes | 2533 | F | <i>Cobitis elongata</i> | N | Y | | U1 | 0 | 3 | 3 | 1 | 12 | | |
| Rivers & lakes | 1107 | F | <i>Salmo marmoratus</i> | N | Y | | U1 | 0 | 2 | 2 | 3 | 10 | | |
| Rivers & lakes | 4064 | I | <i>Theodoxus transversalis</i> | N | Y | Y | XX | 0 | 3 | 3 | 0 | 9 | | X |
| Rivers & lakes | 1152 | F | <i>Aphanius fasciatus</i> | N | Y | | XX | 0 | 2 | 2 | 2 | 8 | | |
| Rivers & lakes | 1137 | F | <i>Barbus plebejus</i> | N | Y | Y | U1 | 0 | 2 | 2 | 2 | 8 | | |
| Rivers & lakes | 1097 | F | <i>Lethenteron zanandreai</i> | N | Y | Y | U1 | 0 | 2 | 2 | 2 | 8 | | |
| Rivers & lakes | 5088 | F | <i>Barbus cyclolepis</i> | N | Y | Y | U1 | 0 | 2 | 2 | 2 | 8 | | |
| Rivers & lakes | 1120 | F | <i>Alburnus albidus</i> | N | Y | | XX | 0 | 2 | 2 | 1 | 6 | | |
| Rivers & lakes | 5289 | F | <i>Chalcalburnus chalcooides</i> | N | Y | | XX | 0 | 3 | 2 | 0 | 6 | X | X |
| Rivers & lakes | 4046 | I | <i>Cordulegaster heros</i> | N | Y | Y | FV | 0 | 3 | 2 | 0 | 6 | | |
| Rivers & lakes | 1139 | F | <i>Rutilus frisii meidingeri</i> | N | Y | Y | XX | 0 | 2 | 2 | 0 | 4 | | |
| Rivers & lakes | 1136 | F | <i>Rutilus rubilio</i> | N | Y | | U2 | 1 | 2 | 1 | 1 | 4 | | |
| Sparsely vegetated land | 7150 | H | Depressions on peat substrates of the Rhynchosporion | N | Y | | U2 | 0 | 10 | 14 | 3 | 170 | | |
| | | | | | | | | | | | | | | |
| Sparsely vegetated land | 6110 | H | Rupicolous calcareous or basophilic grasslands of the Alysso-Sedion albi | Y | Y | | U1 | 0 | 12 | 12 | 2 | 168 | X | |
| | | | | | | | | | | | | | | |
| Sparsely vegetated land | 8220 | H | Siliceous rocky slopes with chasmophytic vegetation | N | Y | | FV | 0 | 13 | 10 | 1 | 143 | X | |
| | | | | | | | | | | | | | | |
| Sparsely vegetated land | 8310 | H | Caves not open to the public | N | Y | | U2 | 2 | 12 | 8 | 2 | 120 | X | X |
| | | | | | | | | | | | | | | |
| Sparsely vegetated land | 8230 | H | Siliceous rock with pioneer vegetation of the Sedo-Scleranthion or of the Sedo albi-Veronicion dillemii | N | Y | | U1 | 0 | 10 | 10 | 1 | 110 | X | X |
| | | | | | | | | | | | | | | |
| Sparsely vegetated land | 8210 | H | Calcareous rocky slopes with chasmophytic vegetation | N | Y | | FV | 1 | 11 | 9 | 1 | 110 | X | |
| | | | | | | | | | | | | | | |
| Sparsely vegetated land | 8160 | H | Medio-European calcareous scree of hill and montane levels | Y | Y | | FV | 1 | 9 | 7 | 1 | 72 | | |
| | | | | | | | | | | | | | | |
| Sparsely vegetated land | 8150 | H | Medio-European upland siliceous scree | N | Y | | FV | 0 | 7 | 8 | 1 | 63 | | |
| | | | | | | | | | | | | | | |
| Sparsely vegetated land | 1210 | H | Annual vegetation of drift lines | N | Y | | U1 | 0 | 6 | 5 | 2 | 42 | X | X |
| | | | | | | | | | | | | | | |

| | | | | | | | | | | | | | | | |
|--------------------------|------|---|--|---|---|---|--|-----------|---|----|----|----|-----|---|---|
| | | | | | | | | U1 | 0 | 5 | 5 | 2 | 35 | X | X |
| Sparingly vegetated land | 2110 | H | Embryonic shifting dunes | N | Y | | | FV | 0 | 4 | 2 | 1 | 12 | | |
| Sparingly vegetated land | 1230 | H | Vegetated sea cliffs of the Atlantic and Baltic Coasts | N | Y | | | FV | 0 | | | | | | |
| Sparingly vegetated land | 8110 | H | Siliceous scree of the montane to snow levels (<i>Androsacetalia alpinae</i> and <i>Galeopsietalia ladani</i>) | N | Y | | | FV | 0 | 6 | 2 | 0 | 12 | | |
| Sparingly vegetated land | 8120 | H | Calcareous and calcshist screes of the montane to alpine levels (<i>Thlaspietea rotundifolii</i>) | N | Y | | | FV | 0 | 4 | 3 | 0 | 12 | | |
| Sparingly vegetated land | 1220 | H | Perennial vegetation of stony banks | N | Y | | | U1 | 0 | 3 | 3 | 0 | 9 | | |
| Sparingly vegetated land | 8240 | H | Limestone pavements | Y | Y | | | XX | 0 | 3 | 1 | 0 | 3 | | |
| Sparingly vegetated land | 1240 | H | Vegetated sea cliffs of the Mediterranean coasts with endemic <i>Limonium</i> spp. | N | Y | | | FV | 0 | 2 | 0 | 0 | 0 | X | |
| Sparingly vegetated land | 8130 | H | Western Mediterranean and thermophilous scree | N | Y | | | FV | 0 | 2 | 0 | 0 | 0 | | |
| Sparingly vegetated land | 1058 | I | <i>Maculinea arion</i> | N | | Y | | U2 | 0 | 12 | 21 | 16 | 444 | X | |
| Sparingly vegetated land | 1283 | R | <i>Coronella austriaca</i> | N | | Y | | U1 | 0 | 11 | 11 | 11 | 242 | X | |
| Sparingly vegetated land | 1261 | R | <i>Lacerta agilis</i> | N | | Y | | U1 | 0 | 11 | 12 | 9 | 231 | X | |
| Sparingly vegetated land | 1329 | M | <i>Plecotus austriacus</i> | N | | Y | | XX | 0 | 10 | 10 | 5 | 150 | | |
| Sparingly vegetated land | 1057 | I | <i>Parnassius apollo</i> | N | | Y | | U2 | 0 | 7 | 13 | 7 | 140 | | |
| Sparingly vegetated land | 1332 | M | <i>Vespertilio murinus</i> | N | | Y | | XX | 3 | 11 | 10 | 0 | 110 | | |
| Sparingly vegetated land | 1256 | R | <i>Podarcis muralis</i> | N | | Y | | FV | 2 | 9 | 5 | 5 | 90 | X | |
| Sparingly vegetated land | 1379 | P | <i>Mannia triandra</i> | N | | Y | | U1 | 0 | 5 | 6 | 4 | 50 | | |
| Sparingly vegetated land | 4066 | P | <i>Asplenium adulterinum</i> | N | Y | Y | | U1 | 0 | 6 | 5 | 2 | 42 | | |
| Sparingly vegetated land | 1421 | P | <i>Trichomanes speciosum</i> | N | Y | Y | | FV | 1 | 6 | 4 | 1 | 30 | | |
| Sparingly vegetated land | 5365 | M | <i>Pipistrellus savii</i> | N | | Y | | XX | 1 | 5 | 4 | 0 | 20 | X | |
| Sparingly vegetated land | 2016 | M | <i>Pipistrellus kuhlii</i> | N | | Y | | FV | 2 | 6 | 3 | 0 | 18 | X | |
| Sparingly vegetated land | 1289 | R | <i>Telescopus falax</i> | N | | Y | | U2 | 0 | 3 | 4 | 1 | 15 | | |
| Sparingly vegetated land | 1284 | R | <i>Coluber viridiflavus</i> | N | | Y | | U1 | 0 | 3 | 2 | 1 | 9 | | |
| Sparingly vegetated land | 1241 | R | <i>Podarcis melisellensis</i> | N | | Y | | U1 | 0 | 2 | 2 | 2 | 8 | | |
| Sparingly vegetated land | 1250 | R | <i>Podarcis sicula</i> | N | | Y | | XX | 0 | 2 | 2 | 2 | 8 | | |
| Sparingly vegetated land | 1295 | R | <i>Vipera ammodytes</i> | N | | Y | | U1 | 0 | 2 | 2 | 2 | 8 | X | |

| | | | | | | | | | | | | | | | |
|-------------------------|------|---|------------------------------|---|--|---|---|----|---|---|---|---|---|--|--|
| Sparsely vegetated land | 1186 | A | <i>Proteus anguinus</i> | Y | | Y | Y | U1 | 0 | 2 | 1 | 2 | 6 | | |
| Sparsely vegetated land | 1333 | M | <i>Tadarida teniotis</i> | N | | Y | | XX | 0 | 3 | 2 | 0 | 6 | | |
| Sparsely vegetated land | 1493 | P | <i>Sisymbrium supinum</i> | N | | Y | Y | U2 | 0 | 2 | 2 | 0 | 4 | | |
| Sparsely vegetated land | 1458 | P | <i>Moehringia tommasinii</i> | N | | Y | Y | U1 | 0 | 2 | 1 | 0 | 2 | | |

8. Appendix 2. List of species and habitats types of the Pannonian region

- Different colours are used for different habitat groups. Species/habitat types present in only one MS are included.
- H= habitat type, A = amphibian, I = invertebrate, M = mammal, P = plant, R = reptile
- **Prio** = priority habitat type or species
- **I, II and IV** refer to Annexes of the Habitats Directive.
- **PAN CS** = Conservation status at the Pannonian region. Red = unfavourable-bad, amber = unfavourable-inadequate, grey = unknown.
- **Positive trend**: positive trend for population & habitat for species or area of the habitat were used. 0 means that there was not any positive trend reported under the used parameters.
- **Criterion A**= number of MS where species/habitat type is present, **Criterion B** = species/habitat types at U2, U1 or unknown status and **Criterion C**= negative trend and **A(B+C)** = the agreed algorithm.
- Cells in yellow highlight the data that was used for ranking the habitat types and species.

| Habitat group | N2K code | Taxonomical group | Description | Prio | I | II | IV | V | PAN CS | Positive trends | Criterion A | Criterion B | Criterion C | A*(B+C) | Occur in BLS | Occur in STE |
|---------------|----------|-------------------|--|------|---|----|----|---|--------|-----------------|-------------|-------------|-------------|---------|--------------|--------------|
| Forests | 91F0 | H | Riparian mixed forests of Quercus robur, Ulmus laevis and Ulmus minor, Fraxinus excelsior or Fraxinus angustifolia, along the great rivers (Ulmenerum minoris) | N | Y | | | | U2 | 0 | 4 | 7 | 2 | 36 | X | X |
| Forests | 91I0 | H | Euro-Siberian steppic woods with Quercus spp. | Y | Y | | | | U2 | 0 | 4 | 7 | 2 | 36 | X | X |
| Forests | 91E0 | H | Alluvial forests with Alnus glutinosa and Fraxinus excelsior (Alno-Padion, Alnion incanae, Salicion albae) | Y | Y | | | | U2 | 0 | 3 | 6 | 2 | 24 | X | |
| Forests | 9180 | H | Tilio-Acerion forests of slopes, scree and ravines | Y | Y | | | | U2 | 0 | 3 | 5 | 1 | 18 | X | |

| | | | | | | | | | | | | | | | | |
|---------|------|---|--|---|---|---|---|--|----|---|---|---|---|----|---|---|
| Forests | 91M0 | H | Pannonian-Balkanic turkey oak – sessile oak forests | N | Y | | | | U2 | 0 | 3 | 4 | 2 | 18 | X | X |
| Forests | 91G0 | H | Pannonic woods with <i>Quercus petraea</i> and <i>Carpinus betulus</i> | Y | Y | | | | U2 | 1 | 3 | 5 | 0 | 15 | X | |
| Forests | 91H0 | H | Pannonic woods with <i>Quercus pubescens</i> | Y | Y | | | | U2 | 1 | 3 | 4 | 1 | 15 | X | |
| Forests | 91N0 | H | Pannonic inland sand dune thicket (<i>Juniperus-Populetum albae</i>) | Y | Y | | | | U2 | 1 | 2 | 4 | 1 | 10 | | |
| Forests | 9110 | H | Luzulo-Fagetum beech forests | N | Y | | | | U2 | 1 | 2 | 3 | 0 | 6 | | |
| Forests | 9190 | H | Old acidophilous oak woods with <i>Quercus robur</i> on sandy plains | N | Y | | | | U1 | 1 | 2 | 3 | 0 | 6 | | |
| Forests | 9130 | H | Asperulo-Fagetum beech forests | N | Y | | | | U2 | 0 | 2 | 2 | 0 | 4 | | |
| Forests | 9150 | H | Medio-European limestone beech forests of the Cephalanthero-Fagion | N | Y | | | | U2 | 0 | 2 | 2 | 0 | 4 | | |
| Forests | 1326 | M | <i>Plecotus auritus</i> | N | | Y | | | U1 | 0 | 3 | 4 | 4 | 24 | | |
| Forests | 1188 | A | <i>Bombina bombina</i> | N | | Y | Y | | U1 | 0 | 4 | 4 | 2 | 24 | X | X |
| Forests | 1088 | I | <i>Cerambyx cerdo</i> | N | | Y | Y | | U1 | 0 | 4 | 5 | 1 | 24 | X | X |
| Forests | 1323 | M | <i>Myotis bechsteinii</i> | N | | Y | Y | | XX | 0 | 4 | 4 | 2 | 24 | X | |
| Forests | 1304 | M | <i>Rhinolophus ferrumequinum</i> | N | | Y | Y | | U1 | 1 | 4 | 5 | 1 | 24 | X | X |
| Forests | 1993 | A | <i>Triturus dobrogicus</i> | N | | Y | | | U1 | 2 | 4 | 5 | 1 | 24 | X | X |
| Forests | 1214 | A | <i>Rana arvalis</i> | N | | Y | | | U1 | 0 | 3 | 4 | 3 | 21 | | |
| Forests | 1207 | A | <i>Rana lessonae</i> | N | | Y | | | U1 | 0 | 3 | 4 | 3 | 21 | | |
| Forests | 1193 | A | <i>Bombina variegata</i> | N | | Y | Y | | U1 | 0 | 4 | 4 | 1 | 20 | X | |
| Forests | 1308 | M | <i>Barbastella barbastellus</i> | N | | Y | Y | | XX | 0 | 3 | 3 | 2 | 15 | X | |
| Forests | 1320 | M | <i>Myotis brandtii</i> | N | | Y | | | XX | 1 | 3 | 3 | 2 | 15 | | |
| Forests | 1322 | M | <i>Myotis nattereri</i> | N | | Y | | | U1 | 0 | 3 | 3 | 2 | 15 | | |
| Forests | 1331 | M | <i>Nyctalus leisleri</i> | N | | Y | | | XX | 0 | 3 | 3 | 2 | 15 | | |
| Forests | 4052 | I | <i>Odontopodisma rubripes</i> | N | | Y | Y | | XX | 0 | 3 | 3 | 2 | 15 | | |
| Forests | 1902 | P | <i>Cypripedium calceolus</i> | N | | Y | Y | | U1 | 1 | 3 | 4 | 0 | 12 | | |
| Forests | 1363 | M | <i>Felis silvestris</i> | N | | Y | | | U2 | 0 | 2 | 3 | 3 | 12 | X | |
| Forests | 1084 | I | <i>Osmoderma eremita</i> | Y | | Y | | | XX | 0 | 3 | 4 | 0 | 12 | X | X |
| Forests | 1209 | A | <i>Rana dalmatina</i> | N | | Y | | | U1 | 1 | 3 | 3 | 1 | 12 | X | |
| Forests | 4068 | P | <i>Adenophora liliifolia</i> | N | | Y | Y | | U2 | 0 | 3 | 3 | 1 | 12 | | |
| Forests | 1281 | R | <i>Elaphe longissima</i> | N | | Y | | | XX | 2 | 3 | 3 | 0 | 9 | X | |

| | | | | | | | | | | | | | | | |
|------------|------|---|---|---|---|---|--|----|---|---|---|---|----|---|---|
| Forests | 1341 | M | <i>Muscardinus avellanarius</i> | N | | Y | | XX | 0 | 3 | 3 | 0 | 9 | X | |
| Forests | 1314 | M | <i>Myotis daubentonii</i> | N | | Y | | XX | 3 | 3 | 2 | 1 | 9 | X | |
| Forests | 1052 | I | <i>Hypodryas maturna</i> | N | | Y | | U1 | 0 | 3 | 2 | 1 | 9 | | X |
| Forests | 1166 | A | <i>Triturus cristatus</i> | N | | Y | | U1 | 0 | 3 | 3 | 0 | 9 | | |
| Forests | 1342 | M | <i>Dryomys nitedula</i> | N | | Y | | U2 | 0 | 2 | 3 | 1 | 8 | X | |
| Forests | 5037 | R | <i>Lacerta vivipara pannonica</i> | N | | Y | | U1 | 0 | 2 | 2 | 2 | 8 | | |
| Forests | 4036 | I | <i>Leptidea morsei</i> | N | | Y | | U2 | 0 | 2 | 3 | 1 | 8 | X | |
| Forests | 1328 | M | <i>Nyctalus lasiopterus</i> | N | | Y | | U1 | 0 | 2 | 2 | 2 | 8 | | |
| Forests | 1167 | A | <i>Triturus carnifex</i> | N | | Y | | U2 | 0 | 2 | 3 | 1 | 8 | | |
| Forests | 1083 | I | <i>Lucanus cervus</i> | N | | Y | | FV | 0 | 4 | 2 | 0 | 8 | X | X |
| Forests | 1086 | I | <i>Cucujus cinnaberinus</i> | N | | Y | | XX | 1 | 3 | 2 | 0 | 6 | | |
| Forests | 1079 | I | <i>Limoniscus violaceus</i> | N | | Y | | U2 | 0 | 2 | 3 | 0 | 6 | | |
| Forests | 1361 | M | <i>Lynx lynx</i> | N | | Y | | U2 | 0 | 2 | 3 | 0 | 6 | X | |
| Forests | 5003 | M | <i>Myotis alcathoe</i> | N | | Y | | U1 | 1 | 2 | 2 | 1 | 6 | X | |
| Forests | 1324 | M | <i>Myotis myotis</i> | N | | Y | | U1 | 3 | 3 | 2 | 0 | 6 | X | X |
| Forests | 1312 | M | <i>Nyctalus noctula</i> | N | | Y | | FV | 0 | 3 | 2 | 0 | 6 | X | |
| Forests | 1317 | M | <i>Pipistrellus nathusii</i> | N | | Y | | FV | 1 | 3 | 2 | 0 | 6 | X | |
| Forests | 5009 | M | <i>Pipistrellus pygmaeus</i> | N | | Y | | XX | 2 | 3 | 2 | 0 | 6 | X | |
| Forests | 5365 | M | <i>Pipistrellus savii</i> | N | | Y | | FV | 2 | 3 | 2 | 0 | 6 | X | |
| Forests | 4032 | I | <i>Dioszeghyana schmidtii</i> | N | | Y | | U1 | 1 | 3 | 2 | 0 | 6 | | |
| Forests | 1352 | M | <i>Canis lupus</i> | Y | | Y | | U1 | 2 | 2 | 2 | 0 | 4 | X | X |
| Forests | 1305 | M | <i>Rhinolophus euryale</i> | N | | Y | | U1 | 2 | 2 | 1 | 1 | 4 | X | |
| Forests | 4026 | I | <i>Rhynsodes sulcatus</i> | N | | Y | | XX | 0 | 2 | 2 | 0 | 4 | | |
| Forests | 4027 | I | <i>Arytrura musculus</i> | N | | Y | | U1 | 0 | 2 | 2 | 0 | 4 | | X |
| Forests | 4057 | I | <i>Chilostoma banaticum</i> | N | | Y | | U1 | 2 | 2 | 2 | 0 | 4 | | |
| Forests | 1078 | I | <i>Callimorpha quadripunctaria</i> | Y | | Y | | FV | 1 | 4 | 1 | 0 | 4 | X | X |
| Forests | 1309 | M | <i>Pipistrellus pipistrellus</i> | N | | Y | | FV | 1 | 3 | 1 | 0 | 3 | X | |
| Forests | 1332 | M | <i>Vespertilio murinus</i> | N | | Y | | XX | 3 | 3 | 1 | 0 | 3 | | |
| Forests | 1087 | I | <i>Rosalia alpina</i> | Y | | Y | | XX | 2 | 2 | 1 | 0 | 2 | X | |
| Grasslands | 6260 | H | Pannonic sand steppes | Y | Y | | | U2 | 0 | 4 | 7 | 4 | 44 | X | X |
| Grasslands | 6250 | H | Pannonic loess steppic grasslands | Y | Y | | | U2 | 0 | 4 | 6 | 3 | 36 | | |
| Grasslands | 6410 | H | Molinia meadows on calcareous, peaty or clayey-silt-laden soils (<i>Molinion caeruleae</i>) | N | Y | | | U2 | 0 | 4 | 6 | 2 | 32 | X | X |

| | | | | | | | | | | | | | | | | |
|------------|------|---|--|---|---|---|---|--|----|---|---|---|---|----|---|---|
| Grasslands | 6440 | H | Alluvial meadows of river valleys of the <i>Cnidion dubii</i> | N | Y | | | | U2 | 0 | 4 | 6 | 2 | 32 | X | X |
| Grasslands | 1530 | H | Pannonic salt steppes and salt marshes | Y | Y | | | | U2 | 0 | 4 | 5 | 2 | 28 | X | X |
| Grasslands | 6430 | H | Hydrophilous tall herb fringe communities of plains and of the montane to alpine levels | N | Y | | | | U2 | 0 | 4 | 6 | 1 | 28 | X | X |
| Grasslands | 6510 | H | Lowland hay meadows (<i>Alopecurus pratensis</i> , <i>Sanguisorba officinalis</i>) | N | Y | | | | U2 | 1 | 4 | 6 | 1 | 28 | X | X |
| Grasslands | 6240 | H | Sub-Pannonic steppic grasslands | Y | Y | | | | U2 | 0 | 3 | 5 | 3 | 24 | X | |
| Grasslands | 6210 | H | Semi-natural dry grasslands and scrubland facies on calcareous substrates (<i>Festuco-Brometalia</i>) (* important orchid sites) | N | Y | | | | U2 | 0 | 3 | 5 | 2 | 21 | X | |
| Grasslands | 2340 | H | Pannonic inland dunes | Y | Y | | | | U2 | 0 | 3 | 4 | 1 | 15 | | |
| Grasslands | 1340 | H | Inland salt meadows | Y | Y | | | | U2 | 0 | 2 | 4 | 2 | 12 | | |
| Grasslands | 6110 | H | Rupicolous calcareous or basophilic grasslands of the <i>Alyssum-Sedion albi</i> | Y | Y | | | | XX | 0 | 3 | 4 | 0 | 12 | X | |
| Grasslands | 6190 | H | Rupicolous pannonic grasslands (<i>Stipo-Festucetalia pallentis</i>) | N | Y | | | | U1 | 0 | 3 | 2 | 1 | 9 | | |
| Grasslands | 1335 | M | <i>Spermophilus citellus</i> | N | | Y | Y | | U1 | 0 | 4 | 6 | 5 | 44 | X | X |
| Grasslands | 2633 | M | <i>Mustela eversmannii</i> | N | | Y | Y | | XX | 0 | 4 | 5 | 2 | 28 | X | X |
| Grasslands | 4110 | P | <i>Pulsatilla pratensis</i> ssp. <i>hungarica</i> | Y | | Y | Y | | U2 | 0 | 3 | 5 | 4 | 27 | | |
| Grasslands | 4098 | P | <i>Iris humilis</i> ssp. <i>arenaria</i> | N | | Y | Y | | U1 | 1 | 4 | 4 | 2 | 24 | | |
| Grasslands | 4013 | I | <i>Carabus hungaricus</i> | N | | Y | Y | | FV | 1 | 4 | 5 | 1 | 24 | | |
| Grasslands | 1059 | I | <i>Maculinea teleius</i> | N | | Y | Y | | U1 | 0 | 4 | 4 | 2 | 24 | | |
| Grasslands | 4096 | P | <i>Gladiolus palustris</i> | N | | Y | Y | | U2 | 2 | 3 | 6 | 1 | 21 | | |
| Grasslands | 1061 | I | <i>Maculinea nausithous</i> | N | | Y | Y | | U1 | 0 | 3 | 4 | 3 | 21 | | |
| Grasslands | 2093 | P | <i>Pulsatilla grandis</i> | N | | Y | Y | | U1 | 0 | 3 | 3 | 4 | 21 | | X |
| Grasslands | 1617 | P | <i>Angelica palustris</i> | N | | Y | Y | | U2 | 0 | 3 | 5 | 2 | 21 | | |
| Grasslands | 4011 | I | <i>Bolbelasmus unicornis</i> | N | | Y | Y | | XX | 0 | 3 | 5 | 1 | 18 | X | X |
| Grasslands | 1689 | P | <i>Dracocephalum austriacum</i> | N | | Y | Y | | U1 | 1 | 3 | 4 | 2 | 18 | | |
| Grasslands | 4067 | P | <i>Echium russicum</i> | N | | Y | Y | | FV | 1 | 3 | 4 | 2 | 18 | | X |
| Grasslands | 4030 | I | <i>Colias myrmidone</i> | N | | Y | Y | | U2 | 0 | 2 | 4 | 4 | 16 | | X |
| Grasslands | 1014 | I | <i>Vertigo angustior</i> | N | | Y | | | FV | 0 | 4 | 4 | 0 | 16 | X | |

| | | | | | | | | | | | | | | |
|------------|------|---|-----------------------------|---|---|---|----|---|---|---|---|----|---|---|
| Grasslands | 1308 | M | Barbastella barbastellus | N | Y | Y | XX | 0 | 3 | 3 | 2 | 15 | X | |
| Grasslands | 4091 | P | Crambe tataria | N | Y | Y | U1 | 0 | 3 | 2 | 3 | 15 | X | X |
| Grasslands | 1056 | I | Parnassius mnemosyne | N | | Y | U1 | 0 | 3 | 3 | 2 | 15 | X | |
| Grasslands | 4097 | P | Iris aphylla ssp. hungarica | N | Y | Y | U1 | 0 | 3 | 3 | 2 | 15 | | X |
| Grasslands | 1058 | I | Maculinea arion | N | | Y | U2 | 0 | 2 | 4 | 3 | 14 | X | |
| Grasslands | 4004 | M | Microtus oeconomus mehelyi | Y | Y | Y | U2 | 0 | 2 | 3 | 4 | 14 | | |
| Grasslands | 2285 | P | Colchicum arenarium | N | Y | Y | U1 | 1 | 2 | 3 | 3 | 12 | | |
| Grasslands | 1339 | M | Cricetus cricetus | N | | Y | U1 | 1 | 3 | 2 | 2 | 12 | | |
| Grasslands | 2327 | P | Himantoglossum caprinum | N | Y | Y | U1 | 0 | 2 | 3 | 3 | 12 | X | X |
| Grasslands | 1263 | R | Lacerta viridis | N | | Y | XX | 0 | 3 | 3 | 1 | 12 | X | |
| Grasslands | 1307 | M | Myotis blythii | N | Y | Y | U1 | 1 | 3 | 3 | 1 | 12 | X | X |
| Grasslands | 1197 | A | Pelobates fuscus | N | | Y | U1 | 0 | 3 | 3 | 1 | 12 | X | |
| Grasslands | 4055 | I | Stenobothrus eurasius | N | Y | Y | U1 | 0 | 2 | 2 | 4 | 12 | | |
| Grasslands | 4104 | P | Himantoglossum adriaticum | N | Y | Y | U1 | 0 | 2 | 3 | 2 | 10 | | |
| Grasslands | 2203 | P | Onosma tornensis | Y | Y | Y | U1 | 0 | 2 | 2 | 3 | 10 | | |
| Grasslands | 1281 | R | Elaphe longissima | N | | Y | XX | 2 | 3 | 3 | 0 | 9 | X | |
| Grasslands | 4050 | I | Isophya stysi | N | Y | Y | U1 | 1 | 2 | 2 | 2 | 8 | | |
| Grasslands | 1310 | M | Miniopterus schreibersii | N | Y | Y | U2 | 1 | 2 | 3 | 1 | 8 | X | X |
| Grasslands | 1477 | P | Pulsatilla patens | N | Y | Y | U2 | 0 | 2 | 2 | 2 | 8 | | |
| Grasslands | 4074 | P | Dianthus diutinus | Y | Y | Y | U2 | 0 | 2 | 3 | 1 | 8 | | |
| Grasslands | 1060 | I | Lycaena dispar | N | Y | Y | U1 | 0 | 4 | 2 | 0 | 8 | X | X |
| Grasslands | 5003 | M | Myotis alcathoe | N | | Y | U1 | 1 | 2 | 2 | 1 | 6 | X | |
| Grasslands | 1076 | I | Proserpinus proserpina | N | | Y | XX | 1 | 3 | 2 | 0 | 6 | | |
| Grasslands | 2021 | M | Sicista subtilis | N | Y | Y | XX | 0 | 2 | 3 | 0 | 6 | | X |
| Grasslands | 1053 | I | Zerynthia polyxena | N | | Y | FV | 0 | 3 | 1 | 1 | 6 | X | |
| Grasslands | 4020 | I | Pilemia tigrina | N | Y | Y | U1 | 0 | 2 | 2 | 1 | 6 | | |
| Grasslands | 1276 | R | Ablepharus kitaibelii | N | | Y | U1 | 0 | 2 | 2 | 0 | 4 | X | |
| Grasslands | 4022 | I | Probaticus subrugosus | N | Y | Y | XX | 0 | 2 | 2 | 0 | 4 | X | |
| Grasslands | 4087 | P | Serratula lycopifolia | Y | Y | Y | U1 | 0 | 2 | 2 | 0 | 4 | | |
| Grasslands | 2120 | P | Thlaspi jankae | N | Y | Y | FV | 1 | 2 | 1 | 1 | 4 | | |
| Grasslands | 4035 | I | Gortyna borelii lunata | N | Y | Y | U1 | 2 | 2 | 2 | 0 | 4 | | |
| Grasslands | 1327 | M | Eptesicus serotinus | N | | Y | FV | 1 | 3 | 1 | 0 | 3 | X | |
| Grasslands | 1261 | R | Lacerta agilis | N | | Y | XX | 0 | 3 | 1 | 0 | 3 | X | |
| Grasslands | 1321 | M | Myotis emarginatus | N | Y | Y | FV | 3 | 3 | 1 | 0 | 3 | X | X |

| | | | | | | | | | | | | | | | |
|-----------------|------|---|--|---|---|---|---|----|---|---|---|---|----|---|---|
| Grasslands | 1016 | I | Vertigo moulinsiana | N | | Y | | FV | 0 | 3 | 1 | 0 | 3 | X | |
| Grasslands | 1256 | R | Podarcis muralis | N | | Y | | XX | 0 | 2 | 1 | 0 | 2 | X | |
| Heaths & scrubs | 40A0 | H | Subcontinental peri-Pannonic scrub | Y | Y | | | U2 | 1 | 4 | 6 | 2 | 32 | X | |
| Heaths & scrubs | 4030 | H | European dry heaths | N | Y | | | U2 | 0 | 3 | 4 | 2 | 18 | X | |
| Heaths & scrubs | 5130 | H | Juniperus communis formations on heaths or calcareous grasslands | N | Y | | | U1 | 0 | 3 | 4 | 0 | 12 | | |
| Heaths & scrubs | 1067 | I | Lopinaga achine | N | | Y | | U2 | 0 | 3 | 6 | 6 | 36 | | |
| Heaths & scrubs | 1323 | M | Myotis bechsteinii | N | | Y | Y | XX | 0 | 4 | 4 | 2 | 24 | X | |
| Heaths & scrubs | 1304 | M | Rhinolophus ferrumequinum | N | | Y | Y | U1 | 1 | 4 | 5 | 1 | 24 | X | X |
| Heaths & scrubs | 1074 | I | Eriogaster catax | N | | Y | Y | U1 | 2 | 4 | 4 | 1 | 20 | | X |
| Heaths & scrubs | 4011 | I | Bolbelasmus unicornis | N | | Y | Y | XX | 0 | 3 | 5 | 1 | 18 | X | X |
| Heaths & scrubs | 1050 | I | Saga pedo | N | | Y | | XX | 2 | 3 | 4 | 2 | 18 | | |
| Heaths & scrubs | 1056 | I | Parnassius mnemosyne | N | | Y | | U1 | 0 | 3 | 3 | 2 | 15 | X | |
| Heaths & scrubs | 1283 | R | Coronella austriaca | N | | Y | | XX | 0 | 3 | 3 | 1 | 12 | X | |
| Heaths & scrubs | 1363 | M | Felis silvestris | N | | Y | | U2 | 0 | 2 | 3 | 3 | 12 | X | |
| Heaths & scrubs | 1307 | M | Myotis blythii | N | | Y | Y | U1 | 1 | 3 | 3 | 1 | 12 | X | X |
| Heaths & scrubs | 1303 | M | Rhinolophus hipposideros | N | | Y | Y | XX | 3 | 4 | 3 | 0 | 12 | X | X |
| Heaths & scrubs | 1341 | M | Muscardinus avellanarius | N | | Y | | XX | 0 | 3 | 3 | 0 | 9 | X | |
| Heaths & scrubs | 1342 | M | Dryomys nitedula | N | | Y | | U2 | 0 | 2 | 3 | 1 | 8 | X | |
| Heaths & scrubs | 1361 | M | Lynx lynx | N | | Y | Y | U2 | 0 | 2 | 3 | 0 | 6 | X | |
| Heaths & scrubs | 1076 | I | Proserpinus proserpina | N | | Y | | XX | 1 | 3 | 2 | 0 | 6 | | |
| Heaths & scrubs | 1053 | I | Zerynthia polyxena | N | | Y | | FV | 0 | 3 | 1 | 1 | 6 | X | |
| Heaths & scrubs | 1352 | M | Canis lupus | Y | | Y | | U1 | 2 | 2 | 2 | 0 | 4 | X | X |
| Heaths & scrubs | 4014 | I | Carabus variolosus | N | | Y | Y | XX | 0 | 2 | 2 | 0 | 4 | | |
| Heaths & scrubs | 1078 | I | Callimorpha quadripunctaria | Y | | Y | | FV | 1 | 4 | 1 | 0 | 4 | X | X |
| Mires & bogs | 7230 | H | Alkaline fens | N | Y | | | U2 | 0 | 3 | 5 | 3 | 24 | | |
| Mires & bogs | 7140 | H | Transition mires and quaking bogs | N | Y | | | U2 | 0 | 3 | 3 | 2 | 15 | | |
| Mires & bogs | 4081 | P | Cirsium brachycephalum | N | | Y | Y | FV | 0 | 4 | 4 | 3 | 28 | | |
| Mires & bogs | 1428 | P | Marsilea quadrifolia | N | | Y | Y | U2 | 0 | 3 | 5 | 4 | 27 | | X |
| Mires & bogs | 1188 | A | Bombina bombina | N | | Y | Y | U1 | 0 | 4 | 4 | 2 | 24 | X | X |
| Mires & bogs | 1059 | I | Maculinea teleius | N | | Y | Y | U1 | 0 | 4 | 4 | 2 | 24 | | |
| Mires & bogs | 1993 | A | Triturus dobrogicus | N | | Y | | U1 | 2 | 4 | 5 | 1 | 24 | X | X |
| Mires & bogs | 4096 | P | Gladiolus palustris | N | | Y | Y | U2 | 2 | 3 | 6 | 1 | 21 | | |
| Mires & bogs | 1061 | I | Maculinea nausithous | N | | Y | Y | U1 | 0 | 3 | 4 | 3 | 21 | | |

| | | | | | | | | | | | | | | | | |
|--------------|------|---|-----------------------------|---|--|---|--|----|---|---|---|---|----|---|---|--|
| Mires & bogs | 1214 | A | Rana arvalis | N | | Y | | U1 | 0 | 3 | 4 | 3 | 21 | | | |
| Mires & bogs | 1207 | A | Rana lessonae | N | | Y | | U1 | 0 | 3 | 4 | 3 | 21 | | | |
| Mires & bogs | 1617 | P | Angelica palustris | N | | Y | | U2 | 0 | 3 | 5 | 2 | 21 | | | |
| Mires & bogs | 1193 | A | Bombina variegata | N | | Y | | U1 | 0 | 4 | 4 | 1 | 20 | X | | |
| Mires & bogs | 1725 | P | Lindernia procumbens | N | | Y | | U1 | 2 | 3 | 4 | 2 | 18 | | | |
| Mires & bogs | 1037 | I | Ophiogomphus cecilia | N | | Y | | FV | 2 | 4 | 3 | 1 | 16 | X | X | |
| Mires & bogs | 1014 | I | Vertigo angustior | N | | Y | | FV | 0 | 4 | 4 | 0 | 16 | X | | |
| Mires & bogs | 1330 | M | Myotis mystacinus | N | | Y | | U1 | 1 | 3 | 3 | 2 | 15 | X | | |
| Mires & bogs | 1040 | I | Stylurus flavipes | N | | Y | | FV | 2 | 3 | 4 | 1 | 15 | | | |
| Mires & bogs | 4045 | I | Coenagrion ornatum | N | | Y | | FV | 0 | 3 | 3 | 2 | 15 | X | X | |
| Mires & bogs | 1614 | P | Apium repens | N | | Y | | U1 | 0 | 2 | 3 | 4 | 14 | | | |
| Mires & bogs | 4004 | M | Microtus oeconomus mechelyi | Y | | Y | | U2 | 0 | 2 | 3 | 4 | 14 | | | |
| Mires & bogs | 1201 | A | Bufo viridis | N | | Y | | U1 | 0 | 3 | 3 | 1 | 12 | X | | |
| Mires & bogs | 1203 | A | Hyla arborea | N | | Y | | U1 | 0 | 3 | 3 | 1 | 12 | X | | |
| Mires & bogs | 1292 | R | Natrix tessellata | N | | Y | | XX | 0 | 3 | 3 | 1 | 12 | X | | |
| Mires & bogs | 1197 | A | Pelobates fuscus | N | | Y | | U1 | 0 | 3 | 3 | 1 | 12 | X | | |
| Mires & bogs | 1209 | A | Rana dalmatina | N | | Y | | U1 | 1 | 3 | 3 | 1 | 12 | X | | |
| Mires & bogs | 1303 | M | Rhinolophus hipposideros | N | | Y | | XX | 3 | 4 | 3 | 0 | 12 | X | X | |
| Mires & bogs | 4063 | I | Sadleriana pannonica | N | | Y | | U1 | 0 | 2 | 3 | 2 | 10 | | | |
| Mires & bogs | 1220 | R | Emys orbicularis | N | | Y | | FV | 0 | 3 | 3 | 0 | 9 | X | X | |
| Mires & bogs | 1042 | I | Leucorrhinia pectoralis | N | | Y | | U1 | 1 | 3 | 3 | 0 | 9 | | | |
| Mires & bogs | 1166 | A | Triturus cristatus | N | | Y | | U1 | 0 | 3 | 3 | 0 | 9 | | | |
| Mires & bogs | 5037 | R | Lacerta vivipara pannonica | N | | Y | | U1 | 0 | 2 | 2 | 2 | 8 | | | |
| Mires & bogs | 1903 | P | Liparis loeselii | N | | Y | | U1 | 1 | 2 | 3 | 1 | 8 | | | |
| Mires & bogs | 1167 | A | Triturus carnifex | N | | Y | | U2 | 0 | 2 | 3 | 1 | 8 | | | |
| Mires & bogs | 1355 | M | Lutra lutra | N | | Y | | FV | 2 | 4 | 2 | 0 | 8 | X | X | |
| Mires & bogs | 1060 | I | Lycaena dispar | N | | Y | | U1 | 0 | 4 | 2 | 0 | 8 | X | X | |
| Mires & bogs | 1082 | I | Graphoderus bilineatus | N | | Y | | U2 | 0 | 2 | 3 | 0 | 6 | | X | |
| Mires & bogs | 4064 | I | Theodoxus transversalis | N | | Y | | U1 | 0 | 2 | 2 | 1 | 6 | | X | |
| Mires & bogs | 4014 | I | Carabus variolosus | N | | Y | | XX | 0 | 2 | 2 | 0 | 4 | | | |
| Mires & bogs | 4046 | I | Cordulegaster heros | N | | Y | | FV | 0 | 2 | 2 | 0 | 4 | | | |
| Mires & bogs | 1516 | P | Aldrovanda vesiculosa | N | | Y | | U1 | 0 | 2 | 2 | 0 | 4 | | X | |
| Mires & bogs | 4027 | I | Arytrura musculus | N | | Y | | U1 | 0 | 2 | 2 | 0 | 4 | | X | |
| Mires & bogs | 1016 | I | Vertigo moulinsiana | N | | Y | | FV | 0 | 3 | 1 | 0 | 3 | X | | |

| | | | | | | | | | | | | | | |
|----------------|------|---|--|---|---|---|----|---|---|---|---|----|---|---|
| Mires & bogs | 4018 | I | Duvalius hungaricus | N | Y | Y | FV | 0 | 2 | 0 | 1 | 2 | | |
| Mires & bogs | 1898 | P | Eleocharis carniolica | N | Y | Y | FV | 2 | 2 | 1 | 0 | 2 | X | |
| Mires & bogs | 1337 | M | Castor fiber | N | Y | Y | FV | 5 | 3 | 0 | 0 | 0 | | |
| Rivers & lakes | 3260 | H | Water courses of plain to montane levels with the Ranunculion fluitantis and Callitricho-Batrachion vegetation | N | Y | | U2 | 0 | 4 | 6 | 1 | 28 | X | X |
| Rivers & lakes | 3150 | H | Natural eutrophic lakes with Magnopotamion or Hydrocharition - type vegetation | N | Y | | U1 | 0 | 4 | 5 | 1 | 24 | X | X |
| Rivers & lakes | 3270 | H | Rivers with muddy banks with Chenopodion rubri p.p. and Bidention p.p. vegetation | N | Y | | U1 | 0 | 4 | 6 | 0 | 24 | X | X |
| Rivers & lakes | 3130 | H | Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or of the Isoëto-Nanojuncetea | N | Y | | U1 | 0 | 4 | 5 | 0 | 20 | X | X |
| Rivers & lakes | 3160 | H | Natural dystrophic lakes and ponds | N | Y | | U2 | 0 | 3 | 4 | 2 | 18 | X | X |
| Rivers & lakes | 3140 | H | Hard oligo-mesotrophic waters with benthic vegetation of Chara spp. | N | Y | | XX | 0 | 2 | 2 | 0 | 4 | X | X |
| Rivers & lakes | 1032 | I | Unio crassus | N | Y | Y | FV | 1 | 4 | 5 | 2 | 28 | X | |
| Rivers & lakes | 1428 | P | Marsilea quadrifolia | N | Y | Y | U2 | 0 | 3 | 5 | 4 | 27 | | X |
| Rivers & lakes | 1318 | M | Myotis dasycneme | N | Y | Y | U1 | 1 | 4 | 4 | 1 | 20 | | |
| Rivers & lakes | 2522 | F | Pelecus cultratus | N | Y | | XX | 0 | 4 | 5 | 0 | 20 | X | X |
| Rivers & lakes | 1159 | F | Zingel zingel | N | Y | Y | XX | 0 | 4 | 5 | 0 | 20 | X | X |
| Rivers & lakes | 4056 | I | Anisus vorticulus | N | Y | Y | FV | 0 | 4 | 4 | 0 | 16 | X | X |
| Rivers & lakes | 2555 | F | Gymnocephalus baloni | N | Y | Y | FV | 0 | 4 | 4 | 0 | 16 | X | X |
| Rivers & lakes | 1157 | F | Gymnocephalus schraetzer | N | Y | Y | U1 | 0 | 4 | 4 | 0 | 16 | X | X |
| Rivers & lakes | 1037 | I | Ophiogomphus cecilia | N | Y | Y | FV | 2 | 4 | 3 | 1 | 16 | X | X |
| Rivers & lakes | 1160 | F | Zingel streber | N | Y | | U1 | 0 | 4 | 4 | 0 | 16 | X | X |
| Rivers & lakes | 1320 | M | Myotis brandtii | N | | Y | XX | 1 | 3 | 3 | 2 | 15 | | |
| Rivers & lakes | 1040 | I | Stylurus flavipes | N | | Y | FV | 2 | 3 | 4 | 1 | 15 | | |
| Rivers & lakes | 4045 | I | Coenagrion ornatum | N | | Y | FV | 0 | 3 | 3 | 2 | 15 | X | X |
| Rivers & lakes | 1203 | A | Hyla arborea | N | | Y | U1 | 0 | 3 | 3 | 1 | 12 | X | |
| Rivers & lakes | 1292 | R | Natrix tessellata | N | | Y | XX | 0 | 3 | 3 | 1 | 12 | X | |

| | | | | | | | | | | | | | | |
|-------------------------|------|---|--|---|---|---|----|---|---|---|---|----|---|---|
| Rivers & lakes | 1122 | F | <i>Gobio uranoscopus</i> | N | Y | | XX | 0 | 3 | 3 | 1 | 12 | | |
| Rivers & lakes | 2011 | F | <i>Umbra krameri</i> | N | Y | | U1 | 0 | 3 | 3 | 1 | 12 | X | X |
| Rivers & lakes | 1149 | F | <i>Cobitis taenia</i> | N | Y | | FV | 0 | 4 | 3 | 0 | 12 | X | X |
| Rivers & lakes | 1145 | F | <i>Misgurnus fossilis</i> | N | Y | | FV | 1 | 4 | 3 | 0 | 12 | X | X |
| Rivers & lakes | 4063 | I | <i>Sadleriana pannonica</i> | N | Y | Y | U1 | 0 | 2 | 3 | 2 | 10 | | |
| Rivers & lakes | 1314 | M | <i>Myotis daubentonii</i> | N | | Y | XX | 3 | 3 | 2 | 1 | 9 | X | |
| Rivers & lakes | 1220 | R | <i>Emys orbicularis</i> | N | Y | Y | FV | 0 | 3 | 3 | 0 | 9 | X | X |
| Rivers & lakes | 2511 | F | <i>Gobio kessleri</i> | N | Y | | XX | 0 | 3 | 3 | 0 | 9 | | X |
| Rivers & lakes | 1042 | I | <i>Leucorrhinia pectoralis</i> | N | Y | Y | U1 | 1 | 3 | 3 | 0 | 9 | | |
| Rivers & lakes | 1355 | M | <i>Lutra lutra</i> | N | Y | Y | FV | 2 | 4 | 2 | 0 | 8 | X | X |
| Rivers & lakes | 1134 | F | <i>Rhodeus sericeus amarus</i> | N | Y | | FV | 2 | 4 | 2 | 0 | 8 | X | X |
| Rivers & lakes | 6158 | F | <i>Gobio albipinnatus</i> | N | Y | | FV | 0 | 3 | 2 | 0 | 6 | X | X |
| Rivers & lakes | 1082 | I | <i>Graphoderus bilineatus</i> | N | Y | Y | U2 | 0 | 2 | 3 | 0 | 6 | | X |
| Rivers & lakes | 4064 | I | <i>Theodoxus transversalis</i> | N | Y | Y | U1 | 0 | 2 | 2 | 1 | 6 | | X |
| Rivers & lakes | 1146 | F | <i>Sabanejewia aurata</i> | N | Y | | FV | 1 | 3 | 2 | 0 | 6 | X | X |
| Rivers & lakes | 4046 | I | <i>Cordulegaster heros</i> | N | Y | Y | FV | 0 | 2 | 2 | 0 | 4 | | |
| Rivers & lakes | 1163 | F | <i>Cottus gobio</i> | N | Y | | U1 | 0 | 2 | 2 | 0 | 4 | | |
| Rivers & lakes | 1105 | F | <i>Hucho hucho</i> | N | Y | Y | XX | 0 | 2 | 2 | 0 | 4 | | |
| Rivers & lakes | 5345 | F | <i>Rutilus virgo</i> | N | Y | Y | XX | 0 | 2 | 2 | 0 | 4 | | |
| Rivers & lakes | 1516 | P | <i>Aldrovanda vesiculosa</i> | N | Y | Y | U1 | 0 | 2 | 2 | 0 | 4 | | X |
| Rivers & lakes | 1130 | F | <i>Aspius aspius</i> | N | Y | | FV | 2 | 4 | 1 | 0 | 4 | X | X |
| Rivers & lakes | 1138 | F | <i>Barbus meridionalis</i> | N | Y | Y | FV | 0 | 2 | 1 | 0 | 2 | | |
| Rivers & lakes | 1337 | M | <i>Castor fiber</i> | N | Y | Y | FV | 5 | 3 | 0 | 0 | 0 | | |
| Sparsely vegetated land | 8160 | H | Medio-European calcareous scree of hill and montane levels | Y | Y | | FV | 0 | 3 | 2 | 0 | 6 | | |
| | 8310 | H | Caves not open to the public | N | Y | | U1 | 1 | 3 | 2 | 0 | 6 | X | X |
| Sparsely vegetated land | 8210 | H | Calcareous rocky slopes with chasmophytic vegetation | N | Y | | FV | 0 | 3 | 1 | 0 | 3 | X | |
| | 8220 | H | Siliceous rocky slopes with chasmophytic vegetation | N | Y | | FV | 0 | 3 | 1 | 0 | 3 | X | |
| Sparsely vegetated land | 8230 | H | Siliceous rock with pioneer vegetation of the Sedo-Scleranthion or of the Sedo albi-Veronicion dilleii | N | Y | | FV | 0 | 3 | 1 | 0 | 3 | X | X |
| | 8150 | H | Medio-European upland siliceous | N | Y | | FV | 0 | 2 | 0 | 0 | 0 | | |

| | | | scree | | | | | | | | | | | | |
|-------------------------|------|---|------------------------------|---|---|---|--|----|---|---|---|---|----|---|--|
| Sparsely vegetated land | 4075 | P | <i>Dianthus lumnitzeri</i> | Y | Y | Y | | U2 | 0 | 3 | 4 | 2 | 18 | | |
| Sparsely vegetated land | 1329 | M | <i>Plecotus austriacus</i> | N | | Y | | U1 | 1 | 3 | 4 | 2 | 18 | X | |
| Sparsely vegetated land | 1058 | I | <i>Maculinea arion</i> | N | | Y | | U2 | 0 | 2 | 4 | 3 | 14 | X | |
| Sparsely vegetated land | 1283 | R | <i>Coronella austriaca</i> | N | | Y | | XX | 0 | 3 | 3 | 1 | 12 | X | |
| Sparsely vegetated land | 4055 | I | <i>Stenobothrus eurasius</i> | N | Y | Y | | U1 | 0 | 2 | 2 | 4 | 12 | | |
| Sparsely vegetated land | 5365 | M | <i>Pipistrellus savii</i> | N | | Y | | FV | 2 | 3 | 2 | 0 | 6 | X | |
| Sparsely vegetated land | 1261 | R | <i>Lacerta agilis</i> | N | | Y | | XX | 0 | 3 | 1 | 0 | 3 | X | |
| Sparsely vegetated land | 1332 | M | <i>Vespa velutina</i> | N | | Y | | XX | 3 | 3 | 1 | 0 | 3 | | |
| Sparsely vegetated land | 2016 | M | <i>Pipistrellus kuhlii</i> | N | | Y | | FV | 2 | 2 | 1 | 0 | 2 | X | |
| Sparsely vegetated land | 1256 | R | <i>Podarcis muralis</i> | N | | Y | | XX | 0 | 2 | 1 | 0 | 2 | X | |

9. Appendix 3. List of species and habitats types occurring only in Bulgaria and/or Romania

| N2K_code | Description | Priority | I | II | IV | V | BLS-BG | BLS-RO | STE-RO | CON-BG | CON-RO | PAN-RO | MBLS-BG | MBLS-RO |
|----------|---|----------|---|----|----|---|--------|--------|--------|--------|--------|--------|---------|---------|
| 1160 | Large shallow inlets and bays | N | Y | | | | X | X | | | | | | |
| 1150 | Coastal lagoons | Y | Y | | | | X | X | | | | | | |
| 1530 | Pannonic salt steppes and salt marshes | Y | Y | | | | | | X | | | | | |
| 31A0 | Transylvanian hot-spring lotus beds | Y | Y | | | | | | | | | X | | |
| 40C0 | Ponto-Sarmatic deciduous thickets | Y | Y | | | | | | X | X | X | | | |
| 62C0 | Ponto-Sarmatic steppes | Y | Y | | | | X | | X | X | X | | | |
| 62D0 | Oro-Moesian acidophilous grasslands | N | Y | | | | | | | X | | | | |
| 91M0 | Pannonian-Balkanic turkey oak –sessile oak forests | N | Y | | | | | | | | X | X | | |
| 91S0 | Western Pontic beech forests | Y | Y | | | | X | | | | | | | |
| 91X0 | Dobrogean beech forests | Y | Y | | | | | | X | | | | | |
| 91V0 | Dacian Beech forests (Symphyto-Fagion) | N | Y | | | | | | | | | X | | |
| 91W0 | Moesian beech forests | N | Y | | | | | | | X | | | | |
| 91Y0 | Dacian oak & hornbeam forests | N | Y | | | | | | X | | X | | | |
| 91Z0 | Moesian silver lime woods | N | Y | | | | X | | | X | | | | |
| 91AA | Eastern white oak woods | Y | Y | | | | X | | X | X | X | | | |
| 91BA | Moesian silver fir forests | N | Y | | | | | | | X | | | | |
| 91CA | Rhodopide and Balkan Range Scots pine forests | N | Y | | | | | | | | X | | | |
| 92A0 | Salix alba and Populus alba galleries | N | Y | | | | | | | | | X | | |
| 92C0 | Platanus orientalis and Liquidambar orientalis woods (Platanion orientalis) | N | Y | | | | | | | | X | | | |

| | | | | | | | | | | | | | |
|------|---|---|---|---|---|---|---|---|---|---|---|---|--|
| 92D0 | Southern riparian galleries and thickets (Nerio-Tamaricetea and Securinegio tinctoriae) | N | Y | | | | X | X | X | | | | |
| 9560 | Endemic forests with <i>Juniperus</i> spp. | Y | Y | | | | | | X | | | | |
| 5288 | <i>Alburnus mandrensis</i> | N | | Y | | | X | | | | | | |
| 5291 | <i>Alburnus sarmaticus</i> | N | | Y | | | X | | | | | | |
| 5290 | <i>Alburnus schischkovi</i> | N | | Y | | | X | | | | | | |
| 4125 | <i>Alosa immaculata</i> | N | | Y | | Y | X | | | X | | | |
| 4126 | <i>Alosa maeotica</i> | N | | Y | | Y | | X | X | | X | | |
| 2491 | <i>Alosa pontica</i> | N | | Y | | Y | | | | | X | | |
| 4127 | <i>Alosa tanaica</i> | N | | Y | | Y | X | X | X | | | | |
| 2132 | <i>Astragalus peterfii</i> | N | | Y | Y | | | | | | X | | |
| 5265 | <i>Barbus bergi</i> | N | | Y | | Y | X | | | | X | | |
| 5263 | <i>Barbus strumicae</i> | N | | Y | | Y | | | | X | | | |
| 2236 | <i>Campanula romanica</i> | N | | Y | Y | | | | X | | | | |
| 4070 | <i>Campanula serrata</i> | N | | Y | Y | | | | | | X | | |
| 4080 | <i>Centaurea immanuelis-loewii</i> | N | | Y | Y | | | | | X | | | |
| 2253 | <i>Centaurea jankae</i> | N | | Y | Y | | | | X | X | | | |
| 2255 | <i>Centaurea pontica</i> | N | | Y | Y | | | X | | | | | |
| 1286 | <i>Coluber najadum</i> | N | | | Y | | | | | X | | | |
| 1228 | <i>Cyrtopodium kotschyi</i> | N | | | Y | | X | | | X | | | |
| 4103 | <i>Dactylorhiza kalopissii</i> | N | | Y | Y | | | | | X | | | |
| 1350 | <i>Delphinus delphis</i> | N | | | Y | | X | | | | | | |
| 1293 | <i>Elaphe situla</i> | N | | Y | Y | | X | | | X | | | |
| 1277 | <i>Eryx jaculus</i> | N | | | Y | | | | | X | | | |
| 1845 | <i>Fritillaria gussichiae</i> | N | | | Y | | | | | X | | | |
| 2191 | <i>Galium moldavicum</i> | N | | Y | Y | | | | | | X | | |
| 6158 | <i>Gobio vladykovi</i> | N | | Y | | | | X | X | X | X | X | |
| 5365 | <i>Hypsugo savii</i> | N | | | Y | | X | | | | X | | |
| 1251 | <i>Lacerta trilineata</i> | N | | | Y | | X | | | | X | | |
| 1222 | <i>Mauremys caspica</i> | N | | Y | Y | | X | | | | X | | |
| 2609 | <i>Mesocricetus newtoni</i> | N | | Y | Y | | X | | X | X | | | |
| 2079 | <i>Moehringia jankae</i> | N | | Y | Y | | | | X | X | | | |

| | | | | | | | | | | | | | | |
|------|--------------------------------------|---|--|---|---|--|---|---|---|---|---|---|---|---|
| 1356 | <i>Mustela lutreola</i> | N | | Y | Y | | | X | X | | | | | |
| 2617 | <i>Myomimus roachi</i> | N | | Y | Y | | X | | | X | | | | |
| 4039 | <i>Nymphalis vaualbum</i> | N | | Y | Y | | | | | X | X | X | | |
| 1269 | <i>Ophisaurus apodus</i> | N | | | Y | | X | | | X | | | | |
| 1268 | <i>Ophisops elegans</i> | N | | | Y | | | | | X | | | | |
| 1904 | <i>Ophrys argolica</i> | N | | | Y | | X | | | | | | | |
| 1200 | <i>Pelobates syriacus</i> | N | | | Y | | X | | | X | | | | |
| 1351 | <i>Phocoena phocoena</i> | N | | Y | Y | | | | | | | | X | X |
| 1238 | <i>Podarcis erhardii</i> | N | | | Y | | | | | X | | | | |
| 2125 | <i>Potentilla emilii-poppii</i> | N | | Y | Y | | X | | X | X | | | | |
| 4023 | <i>Propomacrus cypriacus</i> | N | | Y | Y | | X | | | X | | | | |
| 4043 | <i>Pseudophilotes bavius</i> | N | | Y | Y | | | | X | | X | | | |
| 1208 | <i>Rana graeca</i> | N | | | Y | | | | | X | | | | |
| 1306 | <i>Rhinolophus blasii</i> | N | | Y | Y | | X | | | X | X | | | |
| 1302 | <i>Rhinolophus mehelyi</i> | N | | Y | Y | | X | | X | X | X | | | |
| 1998 | <i>Romanichthys valsanicola</i> | N | | Y | Y | | | | | | X | | | |
| 1371 | <i>Rupicapra rupicapra balcanica</i> | N | | Y | Y | | | | | X | | | | |
| 2318 | <i>Stipa danubialis</i> | N | | Y | Y | | | | | | X | | | |
| 1219 | <i>Testudo graeca</i> | N | | Y | Y | | X | X | X | X | | | | |
| 4116 | <i>Tozzia carpathica</i> | N | | Y | Y | | | | | X | | | | |
| 1171 | <i>Triturus karelinii</i> | N | | Y | Y | | X | | | X | | | | |
| 4008 | <i>Triturus vulgaris ampelensis</i> | N | | Y | Y | | | | | | X | | | |
| 2300 | <i>Tulipa hungarica</i> | N | | Y | Y | | | | | | X | | | |
| 1349 | <i>Tursiops truncatus</i> | N | | Y | Y | | | | | | | | X | X |
| 2635 | <i>Vormela peregrina</i> | N | | Y | Y | | X | | X | X | | | | |

